

Review Article

Open Access

Case Series of Common Omental and Mesenteric Pathologies

Dhatri Kallala*, Balaji Varaprasad Mallula, Srinadh Boppana, Santhosh Jitta and Govardhan Das Joel

Department of Radiodiagnosis, Kamineni Academy of Medical Sciences and Research Center, LB Nagar, Hyderabad, Telangana, India

ABSTRACT

Radiological imaging plays a crucial role in the diagnosis and management of a wide range of abdominal pathologies, including those affecting the omentum and mesentery. The omentum and mesentery are important structures in the abdominal cavity, serving a variety of functions, such as providing support to the gastrointestinal tract and facilitating the flow of lymph and blood. Pathologies affecting the omentum and mesentery can result in a wide range of clinical presentations, from mild discomfort to life-threatening complications. Omentum and mesentery are involved in various benign and malignant pathologies like omental infarction, omental torsion, encapsulated fat necrosis, fat saponification due to trauma or pancreatitis, mesenteric panniculitis, epiploic appendagitis, omental nodules and caking in infective and malignant conditions like tuberculosis, peritoneal carcinomatosis and metastasis. Sometimes, there exists a diagnostic dilemma between the benign condition like focal encapsulated fat necrosis and malignant condition like liposarcoma, localized peritoneal carcinomatosis. Therefore, Radiological imaging techniques, such as computed tomography (CT) and magnetic resonance imaging (MRI), are invaluable tools for the diagnosis of omental and mesenteric pathologies. In this context, radiologists play a vital role in the accurate interpretation of imaging studies, enabling timely and appropriate management of these complex and often challenging conditions.

*Corresponding author

Dhatri Kallala, Department of Radiodiagnosis, Kamineni Academy of Medical Sciences and Research Center, LB Nagar, Hyderabad, Telangana, India

Received: February 28 2024; **Accepted:** March 12, 2024; **Published:** June 30, 2024

Keywords: Omentum, Mesentery, Omental Infarction, Omental Torsion, Epiploic appendagitis, Omental Nodules, Omental Caking, Mesenteric Panniculitis, Encapsulated Fat Necrosis

Background

Peritoneum is the largest serous membrane of the human body, formed by parietal peritoneum that lines the abdominal cavity and visceral peritoneum, that envelopes abdominal viscera [1, 2]. Peritoneal reflections like omentum, mesentery and intraabdominal ligaments interconnect the viscera with in the abdominal cavity.

Omentum is a predominantly fat-containing sheet containing nerves, blood vessels, lymph channels, connective tissue and fat, present between stomach and other organs. There is greater and lesser omentum. Greater omentum aka epiploon is formed by double layer of peritoneum (4 layers in total) that hangs down from greater curvature of stomach and 1st part of duodenum for a variable length and folds back to finally insert on the anterosuperior surface of transverse colon [3]. Similarly lesser omentum extends from the lesser curvature of stomach to the ligamentum venosum in the liver.

Mesentery is a double layered peritoneal tissue that connects small bowel, portions of the colon to the posterior abdominal wall. It contains adipose tissue, superior and inferior vessels and branches, lymphatics [4]. Small bowel mesentery, transverse and sigmoid mesocolon correspond to the name of the organ it connects to the posterior abdominal wall.

Peritoneal reflection pathologies are mostly overlooked as they are missed during the 1st line of investigation, ultrasound and have gained interest due to the increased role of computed tomography in the diagnostic imaging.

In this article we describe the imaging findings of different omental and mesenteric pathologies we came across at our institute.




Figure 1A - Schematic diagram showing greater and lesser omentum in sagittal section.

Figure 1B - Schematic diagram showing mesentery in axial section[5].

Omental Infarction

Omental infarction with or without torsion is a well-recognised cause of acute abdominal pain which is rarely diagnosed on ultrasound. It is commonly seen in the right lower abdomen, adjacent to the caecum, inferior to the liver due to higher mobility and greater length on the right side. Omental infarction results due to a kink in the venous channel or by omental torsion itself [6]. The cause of primary omental infarction is usually idiopathic whereas secondary omental infarction is seen post surgery, trauma or inflammation. Patients present with abdominal pain which is not associated with gastrointestinal symptoms or fever [7]. Sudden abdominal pain in runners, marathoners and healthy individuals should raise the suspicion of omental infarction due to low omental blood flow. It is usually self limiting and treated with analgesics conservatively. Auto amputation is one of the

consequences during resolution due to fibrosis and retraction. When encountered, complications like abscess formation necessitate surgical intervention. USG may not be very helpful in diagnosing omental infarction. If seen it is visualized as a focal area of hyperechogenicity in the omentum. CT scan is the imaging modality of choice for omental infarction, which is seen as a focal area of fat stranding with hyperdense peripheral halo.



Figure 2 (A and B): Omental infarction with fat necrosis in a 15 year old patient with left sided abdominal pain. CT contrast axial section (Figure 2A) and CT contrast coronal section (Figure 2B) show focal area of fat stranding anterior to the descending colon just beneath the anterior abdominal wall represented by straight and curved arrows.

Omental Torsion

Omental torsion is one of the rare causes of acute abdominal pain. Twisting of the omentum along the long axis, compromising the vascular supply, results in omental torsion [8]. Primary omental torsion is seen in the absence of any intra-abdominal pathology and the thickened segment of omentum rotates around the proximal fixed point. Etiology of primary omental torsion is not well known and is assumed to be due to omental anomalies like bifid omentum, accessory omentum, anomalous vascular supply and kinking [9]. Secondary omental torsion is seen in patients with pre-existing intra-abdominal pathologies like inguinal hernia, abdominal wall hernias, tumors or cysts, post-surgical wound. Factors increasing the intra-abdominal pressure like coughing, labour, lifting heavy weights, heavy meals, forceful vomiting displace the omentum and predispose to omental torsion [10]. Acute appendicitis is the major clinical differential diagnosis of omental torsion.

If visualized on ultrasound, omental torsion is seen as a complex mass with hyper and hypoechoic areas with or without intraperitoneal free fluid. CT abdomen has high sensitivity for omental torsion and seen as omental haziness with whirling of omentum in concentric pattern around central blood vessel (whirl sign). Recurrent episodes of incomplete omental torsion result in omental ball formation, thickened fibrosed omentum. In most cases, it is self limiting, unless it is huge or secondary to an underlying pathology.




Figure 3 (A, B and C): Omental torsion with infarction in a 36 year old male patient who came with acute abdominal pain. Figure 3A and 3B axial CT contrast images and Figure 3C coronal CT contrast Images show swirling of vessels in an area of fat stranding in the left side of abdomen and in the pelvis representing omental torsion.



Figure 4 (A and B): Omental Torsion in a 30 year old Female Patient. CT Abdomen Plain Axial Images show swirling of vessels in right paramedian location, beneath the rectus abdominis muscle.

Epiploic Appendagitis

Epiploic appendages are benign peritoneal outpouches that arise from the serosal surface of the colon which contain adipose tissue and vessels. Epiploic appendagitis is more commonly seen in the left colon and cecum and can measure upto 5 centimeters in length. The rectum does not contain any epiploic appendages. Inflammation of epiploic appendages is a resultant of torsion of epiploic appendages with vascular occlusion. Patients usually present with acute left sided abdominal pain. On ultrasound it is visualized as a round, noncompressible hyperechoic mass without internal vascularity, surrounded by a subtle hypoechoic line. On Computed tomography it is visualized as a fatty central core abutting the colon wall with surrounding inflammatory changes and a thin hyperdense rim-Hyperattenuating rim sign [11]. Presence or absence of central dot of high attenuation representing thrombosed vessels [12].




Figure 5: Ultrasound abdomen in a 33-year-old female with complaints of left iliac fossa pain since 4 days, showing ill-defined focal oval shaped area of increased echogenicity anterior to the sigmoid colon in left iliac fossa. Patient was advised contrast CT Abdomen for further evaluation




Figure 6 (A, B and C): Axial (6A), Coronal (6B) and Sagittal (6C) sections of CT abdomen with contrast show small oval area inflammatory fat stranding surrounded by a soft tissue ring anterior to the sigmoid colon represented by arrows.

Omental Gist

Gastrointestinal stromal tumours are benign/ malignant mesenchymal tumours of Gastrointestinal tract, most commonly arising from stomach and small intestine [13]. Extra gastrointestinal stromal tumours can arise from mesentery, retroperitoneum, omentum, liver [14]. Omental GIST is not a common entity. Patients usually present with complaints of abdominal pain, distension, fatigue, making the early diagnosis difficult and are diagnosed at larger sizes. Majority cases of omental GIST arise from the greater omentum are large sized (usually >10 cm). Prognosis of extra gastrointestinal stromal tumors is worse compared to gastrointestinal stromal tumors. These are KIT + and KIT/PDGFRA mutation driven mesenchymal neoplasms [15]. On plain radiograph it can be seen as soft tissue opacity displacing the bowel loops in case of large tumors. On ultrasound It is seen as large tumor with heteroechogenecity and central necrosis but it is difficult to identify the organ of origin. Computed tomography is used for definite diagnosis, and visualized as soft tissue density mass with central areas of necrosis, seen as fluid fluid level. Calcifications are uncommon. On contrast administration, mild enhancement of the peripheral non necrotic component is seen. Treatment is usually by En block resection and chemotherapy with Imatinib.




Figure 7 (A and B): omental GIST in a 42-year-old male patient who came with complaints of abdominal distension since 2 months. CT abdomen plain coronal (7A) and sagittal (7B) images show large well defined lobulated intraperitoneal soft tissue density mass along the anterior abdominal wall and mesentery extending from the epigastrium to the pelvis, abutting the hepatic flexure of colon, proximal sigmoid colon. No evidence of bowel dilatation was noted.




Figure 8 (A, B, C and D): arterial phase axial (8A), coronal (8B), Sagittal (8C) and venous phase axial (8C) section images of CT Abdomen with contrast show mild heterogenous enhancement of the lesion with few non enhancing central necrotic areas. Possibility of omental, mesenteric GIST and desmoid tumor were given as diagnosis.




Figure 9A: Gross specimen of the tumour after explorative laparotomy and resection of tumour

Figure 9B: Histopathological image showing spindle shaped cells in fascicles with nuclear palisading – Mesenteric GIST with omental deposits.




Figure 10A and 10B: Showing Positivity for CD117 and Ki67 respectively.

Omental Nodules and Caking

Normal omentum has attenuation similar to fat, therefore any disease process in the omentum makes it evident. When the omentum is affected by an inflammatory/ infectious/ neoplastic disease process, the initial response is by omental thickening, nodules which later progresses to form a mass which is referred to as omental cake [16]. Common malignancies associated with omental caking are ovarian malignancy, pseudomyxoma peritonei, stomach and small bowel carcinomas [17].




Figure 11: Omental caking in a 55 year old female patient with right ovarian malignancy. Ultrasound abdomen image showing thickened hypoechoic omentum in the umbilical region – Suggestive of metastasis.




Figure 12 A: CT Abdomen with contrast arterial phase axial section image showing omental nodules in right hypochondrium.

Figure 12 B: CT Abdomen plain axial section image showing omental caking in epigastric and umbilical region in a 55-year-old

female patient with known history of right ovarian carcinoma.

Mesenteric Panniculitis

Mesenteric panniculitis is a benign and chronic inflammatory condition of unknown origin, involving the small intestine or colonic mesentery [18]. It is classified according to pathological changes (i) Chronic nonspecific inflammation, (ii) fat necrosis and (iii) fibrosis. If inflammation and fat necrosis are predominant, the condition is known as mesenteric panniculitis. When fibrosis and retraction are predominant it is known as retractile mesenteritis [19]. On computed tomography the imaging features are high attenuating mesenteric fat, mass effect on neighbouring bowel loops, presence of lymph nodes, hypoattenuating halo sign surrounding the vessels and lymph nodes, hyperattenuating pseudocapsule [20].




Figure 13 (A and B): Mesenteric panniculitis in a 42 year old female patient. CT abdomen plain axial (13A) and coronal (13B) section images showing inflammatory fat stranding around the mesenteric vessels displacing the bowel loops but not displacing the mesenteric vessels.

Encapsulated Fat Necrosis

Intra-peritoneal encapsulated fat necrosis (IPEN) is a rare condition in which Post traumatic or ischaemic insult results in fat degeneration encapsulated within a fibrous capsule in the peritoneal cavity. On CT scans, IPEN typically appears as well-circumscribed, round or oval-shaped mass that is iso- or hypodense relative to adjacent fat tissue. It can be located anywhere within the peritoneal cavity, but is most commonly found in the omentum or mesentery. The fibrous capsule around the necrotic fat is seen as a thin or thick rim of soft tissue density. In some cases, calcifications may be present within the capsule. On ultrasound, IPEN is seen as a heterogeneous, hypoechoic mass with internal echogenic foci and hyperechoic fibrous capsule, with occasional acoustic shadowing. Mild mass effect on the adjacent organs and enhancement of the fibrous capsule in encapsulated fat necrosis on contrast enhanced computed tomography mimics liposarcoma or peritoneal carcinomatosis [21]. Clinical history of the patient is critical in differentiating it from liposarcoma, as patients with encapsulated fat necrosis have a history of surgery/ trauma. Temporal evolution also helps in differentiating the two entities, as encapsulated fat necrosis decreases in size over time. IPEN is typically a benign condition that does not require treatment unless it is causing symptoms or complications such as bowel obstruction.




Figure 14 (A and B): Encapsulated fat necrosis in a 45-year-old female patient who underwent hysterectomy for fibroid uterus

15 days back. CT abdomen plain axial (14A) and Sagittal (14B) images show well defined encapsulated area of fat necrosis in the right Para median location in the infraumbilical region.

Conclusion

A number of mechanical, inflammatory, infective and neoplastic pathologies affect the omentum and mesentery. A thorough understanding of radiological appearance of these conditions is crucial for accurate diagnosis and effective treatment of various abdominal conditions. In case of fat necrosis mimicking liposarcoma and peritoneal carcinomatosis, prior imaging and clinical history helps in the correct diagnosis of fat necrosis.

References

- Devos H, Goethals L, Belsack D, Brucker Y, Allemeersch GJ, et al (2020) Fat misbehaving in the abdominal cavity: A pictorial essay. *Polish Journal of Radiology* 85: 32-38.
- O'Niell K (2018) The peritoneum. TeachMeAnatomy. <https://teachmeanatomy.info/abdomen/areas/peritoneum/>.
- Adam A, Dixon AK, Gillard JH, Schaefer-Prokop C (2021) Grainger & Allison's Diagnostic Radiology: A textbook of medical imaging. Elsevier <https://uomanara.edu.iq/blog/books/grainger-allisons-diagnostic-radiology-a-textbook-of-medical-imaging/>.
- Editors BD (2018) Mesentery (organ): Definition, function and anatomy. Biology Dictionary. <https://biologydictionary.net/mesentery/>.
- Wikimedia Foundation (2023) Mesentery. Wikipedia <https://en.wikipedia.org/wiki/Mesentery>.
- Rasuli B, Bashir O (2012) Omental infarction. Radiopaedia.org. <https://radiopaedia.org/articles/omental-infarction>.
- Barai KP, Knight BC (2011) Diagnosis and management of idiopathic omental infarction: A case report. *International Journal of Surgery Case Reports* 2: 138-140.
- Neto A, Ebouda F (2015) Omental torsion. Radiopaedia.org. <https://radiopaedia.org/articles/omental-torsion?lang=us>.
- Occhionorelli S, Zese M, Cappellari L, Stano R, Vasquez G (2014) Acute abdomen due to primary omental torsion and infarction. *Case Reports in Surgery* 2014: 1-4.
- Tandon AA, Lim KS (2010) Torsion of the greater omentum: A rare preoperative diagnosis. *Indian Journal of Radiology and Imaging* 20: 294-296.
- Singh AK, Gervais DA, Hahn PF, Sagar P, Mueller PR, et al. (2005) Acute Epiploic appendagitis and its mimics. *RadioGraphics* 25: 1521-1534.
- Neto A, Gaillard F (2008) Epiploic appendagitis. Radiopaedia.org. <https://radiopaedia.org/articles/epiploic-appendagitis?lang=us>.
- Silverstone L, Gaillard F (2009) Gastrointestinal stromal tumour. Radiopaedia.org. <https://radiopaedia.org/articles/gastrointestinal-stromal-tumour-1?lang=us>.
- Fagkrezos D, Touloumis Z, Giannila M, Penlidis C, Papaparaskeva K, et al. (2012) Extra-gastrointestinal stromal tumor of the omentum: A rare case report and review of the literature. *Rare Tumors* 4: 141-144.
- Todoroki T, Sano T, Sakurai S, Segawa A, Saitoh T, et al. (2007) Primary omental gastrointestinal stromal tumor (GIST). *World Journal of Surgical Oncology* 5: 66.
- Mamlouk MD, vanSonnenberg E, Shankar S, Silverman SG (2011) Omental cakes: Unusual aetiologies and CT appearances. *Insights into Imaging* 2: 399-408.
- Saber M, Gaillard F (2008) Omental cake. Radiopaedia.org.

- <https://radiopaedia.org/articles/omental-cake?lang=us>.
18. Ashraf A, Singh G (2008) Sclerosing mesenteritis. Radiopaedia.org. <https://radiopaedia.org/articles/sclerosing-mesenteritis-1?lang=us>
19. McLaughlin PD, Filippone A, Maher MM (2013) The “Misty mesentery”: Mesenteric Panniculitis and its mimics. *American Journal of Roentgenology* 200. <https://www.ajronline.org/doi/10.2214/AJR.12.8493>.
20. Horton KM, Lawler LP, Fishman EK (2003) CT findings in sclerosing mesenteritis (panniculitis): Spectrum of disease. *RadioGraphics* 23: 1561-1567.
21. Kamaya A, Federle MP, Desser TS (2011) Imaging manifestations of abdominal fat necrosis and its mimics. *Radio Graphics* 31: 2021-2034.