

## Research Article

## Open Access

## Comparison Between Peripheral and Central Venous Catheters Regarding Venous Pressure and Complications among Critically Ill Patients

Amnah Mohammed Howthan<sup>1\*</sup>, Mona Mohamed El-Hady<sup>1</sup> and Nahed Ahmed Mersal<sup>2</sup>

<sup>1</sup>Faculty of Nursing, Medical Surgical Nursing department, King Abdul Aziz University, Jeddah, Saudi Arabia; Bachelor Degree of Nursing, Register Nurse, King Saud medical city, Riyadh, Saudi Arabia

<sup>1</sup>Associate Professor in Medical Surgical department, Faculty of Nursing, King Abdul Aziz University, Jeddah, Saudi Arabia

<sup>2</sup>Professor of Medical Surgical Nursing King Abdul-Aziz University, Jeddah, Saudi Arabia

### \*Corresponding author

Amnah Mohammed Howthan, Faculty of Nursing, Medical Surgical Nursing department, King Abdul Aziz University, Jeddah, Saudi Arabia. E-mail: ahowthan@ksmc.med.sa

**Received:** June 01, 2021; **Accepted:** June 07, 2021; **Published:** June 21, 2021

### Introduction

Central venous catheter (CVC) known as standard method for hemodynamic monitoring that plays an essential role in critically ill patient management. After recognizing critical condition, measuring, and evaluating the underlying pathophysiological strategies and receiving suitable therapy [1]. Venous pressure is the pressure present within a blood vessel that can be measured directly via an inserted catheter, and it reflects the venous return to the heart [2]. The catheter tip may influence venous pressure measurement [3]. A venous catheter can be inserted either central or peripheral [4].

A qualified healthcare practitioner, an anesthetist, or other medical practitioners, is inserted CVC, but a nurse does preservation and follows up. CVC is placement within the subclavian, internal or external jugular, femoral, basilic, or axillary veins. CVCs contained a single lumen or multiple lumens. Categorized by Central venous catheters are into short-term CVCs and Long-term CVCs [5-7]. It has CVCs, which are imported inside the UK every year annually are nearly 200,000 at intensive care units [5].

In contrast, approximately peripheral catheters are 150 millions and central venous catheters are used each year within the United Kingdom are five million [8]. CVC it utilized for providing intravenous therapy, medication or vasopressors support drug, total parenteral nutrition, blood & chemotherapy, and treated by hemodialysis, and central venous pressure monitoring [9,10].

CVC procedure has many associated complications that increase morbidity & mortality rate, length of hospital stays, and healthcare cost [1,11,12]. Despite the advent of ultrasound-guided vascular cannulation [12]. Problems related to CVC are classified into two Early. Late complications. **Early complications** are also called Mechanical difficulties that happen within the insertion

time of the central lines such as arterial puncture leading to a hematoma, pneumothorax, dysrhythmia, and **Late complications** like the infection is it a severe complication that occurs when a central line is in situ which can lead to sepsis, shock and death. And Thrombotic complications that include venous thrombosis and pulmonary embolism [10].

Complications associated with CVC occur in nearly 15% of patients, mainly 5-19% are mechanical complications, 5-26% are infectious complications, and 2-26% are thrombotic complications [13,14]. Central line-associated bloodstream infection (CLABSI) calculated accounts for one-third of died patients with an attributable mortality of 12%-25% [8].

Central venous pressure (CVP) is an essential monitored medical framework in ICU. described CVP as the fluid that transmits through pressure calculated within the thoracic close to the right atrium (Atefvahid, Hassani, Jafarian, Doyle, & Ahmadi, 2017). Central venous pressure is usually measured hour in ICU all over the world [15].

Electronically measure CVP can be expressed at millimeters of mercury (mmHg) or manually methods with centimeters (cm) of H<sub>2</sub>O more than atmospheric pressure. The CVP is influenced by numerous conditions, including technical and physiologic factors. The usual range of CVP at a healthy person is 3-6 mmHg [7]. While the targeted endpoint of CVP is 15 mmHg with the patient had undergone invasive mechanical ventilation.

A peripheral venous catheter (PVC) is a catheter placed in the vascular during therapy. It introduced using a needle, such as that used to draw blood. It is the most used type of catheter in medicine, and in most cases, it is inserted PVC in the hand or arm vein [16]. The nursing staff can insert PVC. Applied PVC

for both medication and patients care; this makes it probable to manage intravenous fluid, blood, total parenteral nutrition, Also applied veins access for hemodynamic monitoring [17]. The nurses should help choose and select the right distal vein for PVC [18]. Peripheral venous catheters include various curative purposes; however, they have resulted in infectious complications and non-infectious complications. Infectious complications like; pain, hematoma, phlebitis, and infiltration. Non- Infectious complications as; leakage, extravasation, bleeding, and blockage [19]. the less common bloodstream infection is associated with PVC; it occurs around 0.1% or 0.5 per 1,000 catheter days [20]. PVC complications are predominantly joint to enrollment techniques, neither reflection to the catheter or infusate [19].

Peripheral venous pressure (PVP) examination method is reversed from the CVP examination. PVP is calculated through the joining of PVC with a tube of a transducer with pressure. Measurement of PVP is intrusive in a small amount, puts an impact on price, and has the ability to foretell the CVP [21]. Within the critically ill patients, PVP monitoring and works as an alternative for CVP. So, technical problems related to CVP measurement can be ignored if PVC is used [21].

In critically ill patients, they demand to address the evaluation of fluid volume status in a minimum complex way. Practicing a modern hemodynamic observation method at the stand of evidence and scientific reasoning will open a new gateway to less complicated yet effective critical care to patients [1]. Some studies were conducted among critically ill patients and found the agreement's degree is a high between CVP and PVP [1,22]. Stated the monitoring of PVP could be utilized as a normal, price-impacting as well as a minimum intrusive alternative to monitoring the CVP and with low morbidity complications [23,24].

The Nurses are accountable for an estimate and set the patients before inserting venous catheters (either peripheral or central), nursing care and conservation intravenous catheters, and prohibiting the expansion of complications [26]. Therefore, this study will compare peripheral versus central venous catheter pressure and complications among critically ill patients.

## Methods

This study was a quantitative, one group comparative Quasi-experimental design study. It was carried out on three adult ICUs (trauma, neurosurgery, and medical) at King Saud Medical City (KSMC), a general clinic in Riyadh, Saudi Arabia.

Current research data collection was using a venous assessment sheet from August 2018 to June 2019. Three Ethical Approval

was obtained, one from the Faculty of Nursing and another one was from the Faculty of Medicine, KAU (Reference Number: 240-19). And last one from the research center at KSMC (Register Number with KACST, KSA: H-01-R-053). Purposive sampling technique was used, included ICU with central venous catheter (subclavian or jugular vein), male or female patient between 20 years and 60 years of age and willing to participate. While excluded patients with cardiac disorders and elderly, burn patients, patients with a femoral central line, contraindicated patients to place the peripheral intravenous catheter, and skin infection at the intended insertion site. The researchers developed a venous assessment sheet in English language. It was developed it after reviewing the relevant recent literature. venous assessment sheet is a series of a developed checklist designed for gathering information about patient demographic and clinical data, venous catheter characteristics evaluation, venous pressure monitoring, and venous catheter-related complication observation. After gaining ethical approval. The representation of 60 critically ill patients meets the inclusion criteria from that target population recruited from selected ICUs. The Researcher measured venous pressures from central and peripheral catheters three times per day for three days. The researchers observed late complications for CVC from the time of insertion until removal. For PVC, late complications from the time of insertion until difficulties appear.

## Data Analysis

The data analyzed by using SPSS version 24. ANOVA, t-test and Chi-Square test, Inferential statistics, and bland-Altman plots have been made.

## Results

### Demographic and clinical data

Table 3.1 presents the distribution of patients according to the demographic characteristics and clinical data. As shown in this table study sample consists of 60 patients, most of them were male (77%), and 55% were single. Results represent the mean age from the studied sample (60 patients) was  $38.8 \pm 1.5$  years. Regarding the educational level, results display that less than half of the selection was graduated from high school (41.7%). the sample according to clinical data It recognized that the mean weight value was  $(76.70 \pm 21.6)$  regarding body weight. It observed that the mean height was  $(167.9 \pm 9.6)$ , and the mean body mass index was  $26.9 \pm 6.1$ . Additionally, It was shown from the table that the most studied patients admitted to Surgical ICU (51.7%), and less than half of them (37%) were diagnosed with Polytrauma cases. About the connection with mechanical ventilation, the results show that 95% were mechanically ventilated.

**Table 1: The distribution of the studied sample according to the patient's demographic characteristics and clinical data (No:60)**

Demographic Data	Descriptive Statistics	
	N	%
Age		
20-29	17	28.3
30-39	14	23.3
40-49	13	21.7
50-59	16	26.7
Age (Mean ±SD)	38.8±12	
Gender		
Male	46	76.7
Female	14	23.3
Marital status		
Single	33	55.0
Married	27	45.0
Educational level		
Illiterate	3	5
Secondary school	5	8.3
High school	25	41.7
Diploma	13	21.7
Bachelors	14	23.3
Clinical data		
Weight in KG (Mean ±SD)	76.70 ± 21.6	
Height in CM (Mean ±SD)	167.92 ±9.6	
BMI(Mean ±SD)	26.9 ± 6.1	
Diagnosis - Septic shock	1	
Surgical cases	21	
Respiratory cases	9	
Polytrauma cases	22	
Hematology cases	2	
Drug overdose	3	
Renal cases	2	
Unit - Medical ICU	1	
Surgical ICU	31	
Trauma ICU	28	
MV- Yes	57	
No	3	

### Venous Pressure Monitoring Follows Up

Table 3.2.illustrate a comparison between mean CVP and PVP three times daily for three consecutive days. I statistically significant difference was observed among the mean CVP and PVP on the first day ( $p=0.018$ ) but nonsignificant differences in the second and third days ( $p=0.057, 0.202$ , respectively). It was observed that the mean PVP during the three days were higher than CVP ( $11.5 \pm 2.5$ ,  $11.3 \pm 2.2$ ,  $10.8 \pm 4$  vs.  $10.5 \pm 2.1$ ,  $10.6 \pm 2.04$ ,  $9.9 \pm 3.70$ , respectively).

**Table 2: Comparison between the mean central venous pressure and peripheral venous pressure measurements throughout three consecutive days. (N:60)**

Day	CVP (Mean $\pm$ SD)	PVP (Mean $\pm$ SD)	t-value (P-value)
Day 1			
at insertion time	10.36 $\pm$ 2.61	11.51 $\pm$ 3.36	.039 (.039)
2 <sup>nd</sup> time	10.56 $\pm$ 2.58	11.33 $\pm$ 2.67	.113 (.113)
3 <sup>rd</sup> time	10.75 $\pm$ 2.62	11.90 $\pm$ 2.89	.025 (.025)
Total	10.5 $\pm$ 2.1	11.5 $\pm$ 2.5	2.390(0.018)*
Day 2			
1 <sup>st</sup> time	10.68 $\pm$ 2.38	11.40 $\pm$ 2.69	.125 (.125)
2 <sup>nd</sup> time	10.71 $\pm$ 2.21	11.36 $\pm$ 2.59	.142 (.143)
3 <sup>rd</sup> time	10.53 $\pm$ 2.46	11.41 $\pm$ 2.66	.062 (.062)
Total	10.6 $\pm$ 2.04	11.3 $\pm$ 2.2	1.925 (0.057)
Day 3			
1 <sup>st</sup> time	10.35 $\pm$ 3.77	11.06 $\pm$ 4.36	.338 (.338)
2 <sup>nd</sup> time	10.10 $\pm$ 4.13	11.15 $\pm$ 4.68	.195 (.195)
3 <sup>rd</sup> time	9.41 $\pm$ 4.07	10.36 $\pm$ 4.63	.236(.236)
Total	9.9 $\pm$ 3.70	10.8 $\pm$ 4	1.283(0.202)

CVP: Central venous pressure

PVP: Peripheralvenous pressure

\*statistically significant at  $P \leq 0.05$

Table 3.3 Demonstrate a comparison between CVC and PVC related early and late complications. The results show that there were no early complications from both CVC and PVC. Regarding late complications, the findings demonstrate that catheter occlusion was the latest complications for PVC (33%), followed by Extravasation and Infiltration grade 4 (28.4%). In comparison, the most common late complications from CVC was catheter-related infection (5 %). The bacteriological examination showed that the three CVC had a positive outcome, with two of them being gram-positive bacilli, and one had gram-negative bacilli. At the same time, the bacteriological examination for PVC was negative.

**Table 3: Comparison between the central venous catheter and peripheral venous catheter-related complications. (N:60)**

Complications	Group		z test	P
	PVC	CVC		
	N (%)	N (%)		
Late complications				
Venous Spasm	2(3.3)	0 (0.0)	1.431	0.152
Extravasation and Infiltration grade 4 (leakage)	17(28.4)	0 (0.0)	4.592	0.000
Catheter occlusion	33(55)	0 (0.0)	8.563	0.000*
Accidental removal	7(11.7)	0 (0.0)	2.779	0.005
Catheter-related infection (Microorganism)				
Yes	0 (0.0)	3(5)	1.777	0.075
No	60 (100)	57 (95)		
Bacteriological examination:				
Gram-positive bacilli	0 (0.0)	2 (3.3)	1.431	0.152
Gram-negative bacilli	0 (0.0)	1(1.7)		

Table 3.4: Revels that the correlation between PVP and CVP ( $P < 0.05$ ) was significant a correlation coefficient of ( $r = 0.896$ ) indicates a strong positive correlation.



**Table 4: Correlation between the central venous pressure and peripheral venous pressure (N:60)**

PVP Total		CVP Total
	Pearson Correlation	.896**
	Sig. (2-tailed)	.000
	N	60
**. the correlation is significant at the 0.01 level (2-tailed).		

## Discussion

### Demographic and Medical Data

The recent study performed 60 adult patients of both genders. The results show that the total age of the researched required sample was  $38.8 \pm 1.5$  years. Most participants were males and single. Lower than half of patients graduated from high school. The results present that most patients admitted to surgical ICU were diagnosed with polytrauma and connected to a mechanical ventilator.

### Venous catheter characteristics

In the present investigation, the mean duration of patients on PVC was  $4.1 \pm 0.1$  days. This result is supported by Dao (2017), who report that the average PVC days for the routine replacement group was (4.29 days SD 2.47). Moreover, Randomized control trial demonstrated by, PVC was introduced without complications for a common of 3.73 ( $\pm 2.25$ ) and a more than 10 days in the empirical collection however the catheter of the monitoring team was recorded for 3.28 ( $\pm 1.66$ ) and extreme of week [26].

Additionally, Katuska and colleagues 'feedback displays that a period >4 days was related to diminishing the danger of PVC failed, which concurs with our feedback. A research by Abolfotouh et al. (2014) explained that from the first 24 to 30 hours in all problems were involved ( $P = 0.0001$ ) [27].

The present research demonstrates that the average timing of patients on CVC was  $9.5 \pm 0.8$ . This result has contradicted Hignell and the Infusion Nurses Society. They found that the catheter site is expected in each subclavian or the internal jugular site and needed for greater than two weeks. Furthermore, the present finding illustrates that the CVC removal's most common reason was no massive fluid resuscitation requirement or no indication for the catheter. This result is similar result [28]. They reported that CVC was removed once there was no demand for a massive volume of fluids and damaging intravenous devices (62%). Two central lines were eliminated because of consistent hematoma and thrombosis in the vein. While Infusion Nurses Society stated, the direct elimination of working of CVC is not recommended by gaining temperature [2].

### Venous pressure monitoring

The present study compared the mean PVP and CVP for three consecutive days. The results found a significant comparison within CVP and PVP on the first day but an insignificant difference on the second and third days. It also shows that the mean CVP value was higher than the mean PVP among three days on all readings. In this process the CVP and PVP recorded complicatedly by combination of CVP manometer to the central venous catheter as well as peripheral venous catheter of critically ill patients through the reading of pressure at exact time, three times in four hours of interval [1]. The present investigation observed that the grand mean of PVP for three days is higher than CVP ( $11.28 \pm 2.19$  vs.  $10.38 \pm 2.10$ , respectively). Furthermore, the present finding illustrates that the CVC removal's most common reason was no

massive fluid resuscitation requirement or no indication for the catheter.

In comparison, a last research reported by stated that the PVP and CVP were registered to the closest 1mm Hg at 5 minutes' interval [29]. These similarly conducted a study by [30]. They are stated that different patients' positions may lead to the elbow's flexion and result in an erroneous value in PVP. Also, reported the external compression via the factor or blood pressure cuff and stretching too much in the arm of the catheterized site can obtain the peripheral vein and elevated PVP [31].

### Venous catheter complications

The resulting complications arise from the venous catheter, including early complications and late complications. Regarding early complications, the recent study demonstrated the absence of complications from CVC and PVC. demonstrated that CVC's early complications occur because of variations in numerous factors such as medical expertise, sort of device, and type of method or vessel use. showed that Nurses work in essential way to inhibit the CVC difficulties or catheter-related bloodstream infection; by using standard guidelines like arrangement of an aseptic environment in CVC introduction [31].

A current study detected that few patients have PVC developed venous spasm and absence of venous spasm from CVC. This result consistent with Piperet al.who stated that PVCs usually lead toinfiltration, occlusion, phlebitis or thrombophlebitis, dislodgement, and venous spasm [32,33].

The findings identified that extravasation and infiltration occur for one-third of patients with PVC while not observed at the CVC site. stated that extravasation's incidence andinfiltration of PVC, was 3.5% and 7%, respectively[34]. The PVC complication rate accretion with various things that act as danger like person's age and gender as well as the imbalance veins related to infection maximize the chances of it. announced that Extravasation and Infiltration caused by inappropriate placement of PVC, dislodgement, distal puncture, or erosion linked to relative movement of the patient and the catheter [35].

In the present result, the main reason for PVC removal was catheter occlusion. stated that occlusion can come from mechanical blockage of the PVC's or fibrin deposition on the catheter's tip. It may also phlebitis veins swollen or insertion at a point of flexion, both of which may collapse the catheter and prevent flow [35].

A recent study found accidental removal happened to a few patients with PVC while it did not occur to CVC. The results from the same point of view as Dougherty and Lister. They reported that some peripheral cannulas have wings that help secure the skin device to prevent a piston-like movement of the vein and accidental removal.

### Conclusion and Recommendations

The current study constructs that differences statistically significant between the mean CVP and PVP on the first day. Still, there were no significant differences found on the second and third days. The mean PVP during the three days was higher than CVP was observed. The findings demonstrate that catheter occlusion was the most common late complication for PVC, followed by extravasation and infiltration grade 4. While the most common late complication from CVC was a catheter-related infection. Based on this study's results, the researcher suggests involving PVC as a method for measuring venous pressure in clinical practice

and develop educational programs for healthcare professionals about the care and prevention for PVCs complications [36-67].

## References

1. Dan M A & Varghese LC E M (2015) Correlation between Peripheral Venous Pressure and Central Venous Pressure Monitored by CVP Manometer: an Observational Study. Age (Yrs) 40: 41-60]
2. Gorski L A (2017) The 2016 Infusion Therapy Standards of Practice. Home Healthcare Now, 35: 10-18.
3. Sondergaard S, Parkin G, & Aneman A (2015) CVP: we need to bring clinical use into physiological context. Acta Anaesthesiologica Scandinavica 59: 552-560]
4. Ricard JD, Salomon L, Boyer A, Thierry G, Meybeck A, et al. (2013). Central or Peripheral Catheters for Initial Venous Access of ICU Patients: A Randomized Controlled Trial. Crit Care Med 41: 2108-2115.
5. Smith R N, & Nolan J P (2013) Central venous catheters. BMJ : British Medical Journal, 347, f6570.
6. Tobar JEG, Zuleta ALT (2018) Reviving an Old Technique Forgotten: Insertion Central Venous Catheter by External Jugular Vein: Description of Technical and Series of Cases. Ann Clin Lab Res.2018; Vol.6 No.3:243 DOI: 10.21767/2386-5180.100243.
7. Thomas Hill B (2018) Role of central venous pressure (CVP) monitoring in critical care settings. Nursing Standard 32: 41-48.
8. Becerra M B, Shirley D, & Safdar N (2016) Prevalence, risk factors, and outcomes of idle intravenous catheters: An integrative review. American Journal of Infection Control 44: e167-e172..
9. Luckianow G M, Smith D, Bullen D, & Kaplan L J (2016) Understanding percutaneous and subcutaneous central venous access devices. Journal of the American Academy of PAs, 29: 33-36.
10. Wong A V, Arora N, Olusanya O, Sharif B, Lundin R M, Dhadda A, ... First Intensive Care National Audit Project (ICNAP-1) group (2018). Insertion rates and complications of central lines in the UK population: A pilot study. Journal of the Intensive Care Society 19: 19–25. doi:10.1177/1751143717722914.
11. Kornbau C, Lee K C, Hughes G D, & Firstenberg M S (2015) Central line complications. International journal of critical illness and injury science 5: 170.
12. Bowdle A (2014) Vascular Complications of Central Venous Catheter Placement: Evidence-Based Methods for Prevention and Treatment. J Cardiothorac Vasc Anesth 28: 358-368.
13. Blanco P (2016) Ultrasound-guided vascular cannulation in critical care patients: A practical review. Medicina Intensiva (English Edition) 40: 560-571. doi:10.1016/j.medine.2016.07.002.
14. Kander T, Frigyesi A, Kjedsen-Kragh J, Karlsson H, Rolander F, & Schott U (2013) Bleeding complications after central line insertions: relevance of pre-procedure coagulation tests and institutional transfusion policy. Acta Anaesthesiol Scand 57: 573-579.
15. Magder S (2015) Understanding central venous pressure: not a preload index?. Current opinion in critical care 21: 369-375.
16. Klabunde RE (2014) Cardiovascular Physiology Concepts: CVP. www.cvphysiology.com/Blood%20Pressure/BP020 (accessed: 27 November 2017).
17. Pasalioglu K B, & Kaya H (2014) Catheter indwell time and phlebitis development during peripheral intravenous catheter administration. Pakistan Journal of Medical Sciences, 30(4), 725-730.
18. Arslan M, Yalçın S, Kesik F, Demirci B, & Balçık Ö Ş (2014) Turkish nurses' knowledge about application, care, and complications of peripheral and central venous catheters and port catheters. NERP 4: 11-16.
19. Marsh N, Prac M, Webster J, Larsen E, HlthRes G et al. (2017) Observational Study of Peripheral Intravenous Catheter Outcomes in Adult Hospitalized Patients: A Multivariable Analysis of Peripheral Intravenous Catheter Failure. Journal of Hospital Medicine. 12, PP1-7.
20. Mermel L A (2017) Short-term Peripheral Venous Catheter-Related Bloodstream Infections: A Systematic Review. Clinical Infectious Diseases 65: 1757-1762.
21. Rajeev DS, Sheela Verghese (2017) Peripheral venous pressure: An alternative to central venous pressure? Journal of medical science and clinical research. 83.27 ISSN (e)-2347-176x ISSN (p) 2455-0450 DOI: https://dx.doi.org/10.18535/jmscr/v5i4.154.
22. Johansson E, Hammarskjöld F, Lundberg D, & Arnlind M H (2013) Advantages and disadvantages of PICC compared to other central venous lines: a systematic review of the literature. Acta oncologica 52: 886-892.
23. Prakash J, Rao N S, Kumar S, Raghwendra K H, Saran K et al (2018) Study of Relationship between Central Venous Pressure and Peripheral Venous Pressure during Intraoperative Period in Neurosurgical Patients. Journal of Neuroanaesthesiology and Critical Care 5: 15-20.
24. Sunil R, Vishnu N, & Lakshmi K (2016) Correlation between central venous and peripheral venous pressures in surgical patients. Ain-Shams Journal of Anaesthesiology, 9: 52-56. doi:10.4103/1687-7934.178880.
25. Kumar D, Ahmed S, Ali S, Ray U, Varshney A et al. (2015) Correlation between CVP and PVP with passive leg raise in patients on mechanical ventilation. Indian Journal of Critical Care Medicine 19: 648-654.
26. Johann D A, Danski M T R, Vayego S A, Barbosa D A, Lind J (2016) Risk factors for complications in peripheral intravenous catheters in adults: secondary analysis of a randomized controlled trial. Revistal latino-americana de enfermagem, 24]
27. Miliani K, Taravella R, Thillard D, Chauvin V, Martin E et al. (2017) Peripheral venous catheter-related adverse events: evaluation from a multicentre epidemiological study in France (the CATHEVAL Project). PLoS One, 12(1), e0168637.
28. Korula S, Paul V (2015) Incidence of Complications after Central Venous Cannulation-A Prospective Observational Study. CVP, 2, 4-0.
29. Ravindran R, Komu F, PMA B, Ali A Ramadas K (2017) Peripheral Venous Pressure as a Predictor of Central Venous Pressure during Neurosurgical Procedures. Annals of International Medical and Dental Research 33: 8-42.
30. Radhakrishna N, Singh S, Sharma R, Bajaj V, Taank P (2019) Comparative Study of Venous Pressure Obtained From Central and Peripheral Venous Catheter. International Journal of Biomedical Research, 10(1), e5000.
31. Shah M (2017) "Practice of Nursing Care for Central Venous Catheter Among Icus Nurses in Private Tertiary Care Hospital Peshawar, KP". Published 2017 DOI:10.19080/JOJNHC.02.555585.
32. Piper R, Carr P, Kelsey L, Bulmer A, Keogh S, Doyle B (2018) The mechanistic causes of peripheral intravenous catheter failure based on a parametric computational study. Scientific Reports ; volume 8, Article number: 3441.
33. Kaur p, claire R, Gregory S, Domer and Kevin R Glover (2019) Dangers of peripheral intravenous catheterization : The Forgotten Tourniquet and Other Patient Safety Considerations, Open access peer-reviewed chapter 7. DOI:



- 10.5772/intechopen.83854.
34. Macm M (2017) Peripheral Intravenous Catheter (PIVC) Related Local Complications among Patients in KFCH-Jizan.
35. Kaur p, claire R, Gregory S, Domer and Kevin R, Glover, (2019) Dangers of peripheral intravenous catheterization : The Forgotten Tourniquet and Other Patient Safety Considerations, Open access peer-reviewed chapter 7. DOI: 10.5772/intechopen.83854.
36. Abolfotouh M A, Salam M, Ala'a Bani-Mustafa D W Balkhy H H (2014) Prospective study of incidence and predictors of peripheral intravenous catheter-induced complications. *Therapeutics and clinical risk management*, 10, 993.
37. Adekola O O, Iurhe N K, Raji V A, Desalu I (2018) Central venous catheter insertion in critical illness: Techniques, complications. *Journal of Clinical Sciences*, 15(2), 96.
38. Alexandrou E, Ray-Barruel G, Carr P J, Frost S A, Inwood S et al. (2018) Use of short peripheral intravenous catheters: characteristics, management, and outcomes worldwide *J Hosp Med*. 13(5).
39. Aloush S M, & Alsaraireh F A (2018) Nurses' compliance with central line associated blood stream infection prevention guidelines. *Saudi medical journal* 39: 273–279.
40. Atefvahid P, Hassani K, Jafarian K, Doyle D J, Ahmadi H (2017) Analysis of central venous pressure (CVP) signals using mathematical methods. *Journal Of Clinical Monitoring And Computing* 31: 607-616.
41. Awad SA, El-Soussi AH, Sultan MA, El-Farrash M, Tantawy N M (2011) Effect of Intravenous Catheter Placement on Venous Pressure Reading and the Risk of Complications in Critically Ill Patients.
42. Barb Nickel (2019) Peripheral Intravenous Administration of High-Risk Infusions in Critical Care: A Risk-Benefit Analysis. *Critical Care Nurse* 39: 16-28.
43. Beecham GB, Tackling G (2019) Peripheral Line Placement. In *StatPearls* [Internet]. StatPearls Publishing.
44. Bell T, O'Grady N P (2017) Prevention of Central Line–Associated Bloodstream Infections. *Infectious Disease Clinics* 31: 551-559.
45. Braga L M, Parreira, P M, Oliveira A D S S, Mónico L D S M, Arreguy-Sena C et al. (2018) Phlebitis and infiltration: vascular trauma associated with the peripheral venous catheter. *Revista latino-americana de enfermagem*, 26.
46. Clinical Excellence Commission (2013). Peripheral Intravenous Cannula (PIVC) Insertion and Post Insertion Care in Adult Patients. Guideline, PP1-16.
47. Dao L A T (2017) Comparison of peripherally inserted intravenous catheter complication prevalence: Before and after changing a 96-hour routine replacement standard. California State University, Long Beach.
48. Dibble C T, Kohl B A, Lanken P N (2014) Vascular access issues and Procedures. (Second ed., pp. 104-116.e2) doi:10.1016/B978-1-4160-2455-2.00011-6.
49. Dougherty Lister S (2015) The royal Marsden manual of clinical nursing procedures: 9th edition (14), London: NHS foundation trust. P.862-874.
50. Dunda S E, Demir E, Mefful O J, Grieb G, Bozkurt A et al. (2015) Management, clinical outcomes, complications of acute cannula-related peripheral vein phlebitis of the upper extremity: A retrospective study. *Phlebology* 30: 381-388.
51. Fang S, Yang J, Song L, Jiang Y, Liu Y (2017) Comparison of three types of central venous catheters in patients with malignant tumor receiving chemotherapy. *Patient preference and adherence*, 11, 1197.
52. Hassan N, Hammodi A, Alhubail, R, Rayyan N (2017) Effect of Insertion Site on Risk of Central Line Associated Blood Stream Infection in Critically Ill Patients. *Ann Vasc Med Res*, 4(7), 1081.
53. Helm R E, Klausner J D, Klemperer J D, Flint L M, Huang E (2015) Accepted but unacceptable: peripheral IVC failure. *Journal of Infusion Nursing* 38: 189-203.
54. Hignell, P. (2016). Central Venous Catheters In Adult Patients Self-Learning Module. Vascular Access Clinical Practice Committee, Fraser Health Authority. Version 8, PP1-92.
55. Infusion Nurses Society (2016) Infusion Therapy Standards of Practice. *Journal of Infusion Nursing*, 39(1S).
56. Jahagirdar S, Laxmimani, Athiraman U, Ravishankar M (2013) Migration of subclavian venous catheter tip: Patient positioning in ICU makes a difference. *Indian Journal of Critical Care Medicine* May-June, Vol 17 Issue 3.
57. Lai N M, Chaiyakunapruk N, Lai N A, O'Riordan E, Pau W S C et al. (2016) Catheter impregnation, coating or bonding for reducing central venous catheter-related infections in adults. *Cochrane Database of Systematic Reviews*, (3).
58. Lorente L, Huidobro M S, Martín M M, Jiménez A, Mora M L (2004) Accidental catheter removal in critically ill patients: a prospective and observational study. *Critical Care*, 8(4), R229.
59. Memon J I, Rehmani, R S, Venter J L, Alaithan A, Ahsan I et al. (2010) Central venous catheter practice in an adult intensive care setting in the eastern province of Saudi Arabia. *Saudi Med J* 31: 803-807.
60. Moureau N (2018) Impact and Safety Associated with Accidental Dislodgement of Vascular Access Devices: A Survey of Professions, Settings, and Devices. *Journal of the Association for Vascular Access* 23: 203-215.
61. O'Horan H (2015) Peripheral Venous Cannula Insertion and Management (Adults) Policy Hampshire Hospitals NHS Foundation Trust. PP1-19.
62. Piredda M, Biagioli V, Barrella B, Carpisassi I, Ghinelli R et al. (2016) Factors affecting difficult peripheral intravenous cannulation in adults: a prospective observational study. *Journal of Clinical Nursing* 26: 1074–1084.
63. Rickard CM, Ray-Barruel G (2017) Peripheral intravenous catheter assessment: Beyond phlebitis. *Lancet Haematology* 4: 402-403.
64. Wallis, M. C., McGrail, M., Webster, J., Marsh, N., Gowardman, J., Playford, E. G., & Rickard, C. M. (2014). Risk factors for peripheral intravenous catheter failure: a multivariate analysis of data from a randomized controlled trial. *Infection Control & Hospital Epidemiology* 35: 63-68.
65. Webster J, Osborne S, Rickard CM, New K (2015) Clinically-indicated replacement versus routine replacement of peripheral venous catheters *Cochrane Librarian* 8: CD007798.
66. Zhang Li, Siyu Cao, Nicole Marsh, Gillian Ray-Barruel, Julie Flynn, Emily Larsen, and Claire M. Rickard. "Infection risks associated with peripheral vascular catheters." *Journal of infection prevention* 17, no. 5 (2016): 207-213.
67. Yanagisawa A, Takazawa T, Kanamoto M, Tobe M, Hinohara H et al. (2017) Simultaneous peripheral and central venous pressure monitoring for evaluating cardiac preload in critically ill patients. *Age (years)*, 68, 12.

**Copyright:** ©2021 Amnah Mohammed Howthan, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.