

Research Article

Open Access

Effect of Growth Regulators and NaCl Stress on Callus Formation and Alkaloids Production on Vinca Rosea Plant (Catharanthus Roseus L)

Youssif Mahmoud Ahmed

Department of Biotechnology, Faculty of Agriculture, Al-Azhar Univeristy, Cario, Egypt

ABSTRACT

Vinca alkaloids are a subset of drugs obtained from the periwinkle plant they are naturally extracted from the pink periwinkle plant (*Catharanthus roseus*), and they have been used to treat diabetes, high blood pressure and have been used as disinfectant. The vinca alkaloids are also important for being Cancer resistant there are four major vinca alkaloids in Clinical Use: Vinblastine (VBL), vinorelbine (VRL), vincristine (VCR) and vindesine (VDS). Vinca alkaloids are the second most used class of cancer drugs and will stay among the original cancer therapies. *Catharanthus roseus* is still source used for the powerful antitumor drugs. Callus culture had been done on MS-medium containing Different Concentrations from BA-2, 4-D-NAA-IAA. Where the different effects of growth regulators were studied when different parts of the plant have been cultured from leaves and stems this is to induce callus formation and encourage growth and the concentrations that were prepared: BA (0,1-0, 3-1-2 mg/l)-NAA (0,1-1mg/l)-IAA (0,1 mg/l)-2, 4-D (0,75-1mg/l). And after culture the explants we transferred it to the growth room with a temperature ranging from 27 to 29°C relative humidity around 80% And in complete darkness then we applied the resulting callus tissue under the influence of different concentrations of NaCl as (50-75-100-150mM/L). The results from HPLC analysis for the dry weight of callus were as follows ajmalicine (0.18 mg/g DW), catharanthine (0.07 mg/g DW), serpentine (0.37mg/g DW), vindoline (0.11 mg/g DW).

*Corresponding author

Youssif Mahmoud Ahmed, Department of Biotechnology, Faculty of Agriculture, Al-Azhar Univeristy, Cario, Egypt. Tel: +201093906406; E-mail: youssifmahmoud19999@gmail.com

Received: August 12, 2022; **Accepted:** August 18, 2022; **Published:** August 26, 2022

Keywords: Alkaloids, *Catharanthus Roseus*, Growth Regulators High Performance Liquid Chromatography (Hplc), Fluorescence, Uv

Introduction

It has beautifully full-brimmed leaves and large single flowers with long neck in color Varied between white, pink, purple, and white with a red-eyed cup Five separate sepals with five petals fused and wrapped in a layer or apical turn Pollens with five stamens atop a reciprocal hill with endospermic seed petals, it is an outdoor shrub with spread branches and is used to cover the ground areas between the trees and underneath it, it may rise to 80 cm. So that it seems to you that the flowers remain and do not wither and facilitate the care of them because they tolerate drought and extreme heat, and the most important thing for them is to provide daily irrigation once, especially in the summer and very hot, and it is preferable to water at a time Sunset or shortly before it so that the plant can benefit from the largest amount of water throughout the night, as evaporation decreases and it is a tree that loves shady places, so it is planted under trees or between them, It was found that the pond's eye plant contains more than 50 alkaloids that have a rapid medicinal effect for the treatment of some serious diseases. The aqueous extracts of the eye of the prunus are used as a drink for dentin for gum infections, dental pain and decay, as well as for the dentin of gastrointestinal ulcers, and recently it was possible to know the vital effect of some alkaloid substances in Chronic constipation dentin, More than 40 years ago,

other substances have been shown to have an effective effect in the dentin of many types of cancerous diseases, including: vincristine, vinblastine, Ajmalicine and Serpentine, in addition to the substance Hypertension, which plays a role in the dentin of hypertension. The plants during their growth stages adopt a group Among the metabolites that you use in growth and development, and these materials include alkaloids, evidence, phenols, glycosides, and others. These compounds are important for the plant as they are defensive materials against Pathogens, in addition to their use as pharmaceuticals, food flavors, dyes, perfumes, or pesticides, and plants are considered the continuous source in the work to produce important secondary metabolites, Plant tissue culture is the science of growing plant cells, tissues or organs isolated from mother plant on artificial media. Plant tissue culture is the science of growing plant cells, tissues or organs isolated from mother plant on artificial media [1]. It includes techniques and methods appropriate to research into many botanical disciplines and several practical objectives. Both organized and unorganized growth is possible in vitro [2]. For most of the world's population, medicinal plants are the only source of life-saving medicines. They still serve as an important therapeutic aid in the treatment of human diseases. Early humans were driven to explore their immediate natural surroundings in search of defense mechanisms, longevity, and treatments to relieve pain and discomfort. It led to the use of plants, animal products and other materials [2]. Periwinkle (*Catharanthus roseus*), a member of the family Apocynaceae, highly valued in the horticultural industry. It has beautifully full-brimmed leaves and

large single flowers with long neck in color Varied between white, pink, purple, and white with a red- eyed cup Five separate sepals with five petals fused and wrapped in a layer or apical turn Pollens with five stamens atop a reciprocal hill with endospermic seed petals, it is an outdoor shrub with spread branches and is used to cover the ground areas between the trees and underneath it, it may rise to 80 cm [3]. So that it seems to you that the flowers remain and do not wither and facilitate the care of them because they tolerate drought and extreme heat, and the most important thing for them is to provide daily irrigation once especially in the summer and very hot, and it is preferable to water at a time Sunset or shortly before it so that the plant can benefit from the largest amount of water throughout the night, as evaporation decreases and it is a tree that loves shady places, so it is planted [2]. Catharanthus roseus produces more than 100 monoterpenoids indole alkaloids (TIA) in different organs. Leaves and stems are sources of dark alkaloids, Vincristine and vinblastine are indispensable cancers medicines, while the roots have antihypertensive, ajmalicine and serpentine [4]. Catharanthus roseus (L.) G. Don. (Habit Known as the family Periwinkle Apocynaceae) it has gained commercial importance due to its alkaline contents in various parts of the plant. Some are used in the pharmaceutical industry to treat childhood leukemia, Hodgkin's disease, testicular cancer and carcinoid tumors [3]. Alkaloids are a class of organic compounds consisting of Carbon, hydrogen, nitrogen, and usually oxygen, which are often derived from plants such as C. alkaloids It has physiological effects that make the vinca plant of medicinal value. Some vinca alkaloids are strong muscles Relaxer, others dilate the pupils of the eye [5].

Materials and Methods

The experiments and Cultivation process and chemical analysis were carried out in the Laboratory of Biotechnology Department, Faculty of Agriculture, Al-Azhar University, Nasr City, Cairo, Egypt.

Plant Material

The main plant part used in the cultivation process The parts that were used in cultivation were a stalk 2 cm long [2] and a part of the leaf 1 cm long and 0.5 cm wide free of middle sweat was obtained for Vinca Rosa plant (one years-old produced by explant) cultivated in the farm of Faculty of Agriculture, Al-Azhar University, Nasr City, Cairo, Egypt [3].

Explant Preparation

Surface sterilization of explants The branch of a plant has been collected from garden and it cut into small parts and Separate the leaves from the stem and it cut the stems the plant to small part it is 4 cm long [2] and these parts were put in a big jar, and wash with industrial detergent and they were washed with running water 4 times, and It's surface was sterilized with ethanol at 97% concentration for 2 minutes , Then we washed 3: 5 times with sterile distilled water in a laminar flow hood [2], Now the parts are ready to be cultured [5].

Effect of Growth Regulators on Callus Formation

Sterile uniform explants i.e. 1.0*0.5 cm for leaf blade is free of middle sweat and placed in the outside of it and main stem are a 2 centimeter long placed in italics in media were aseptically transferred to jars containing 30 mL of Morishige and Skoog (MS) medium 2 liters of media were prepared to contain (8.86g/2L MS+ 60g / 2L sucrose + 14 g agar) then we take 1.75 liters from them and were individually transferred into 250 ml jars and different combinations of 2,4-D (2,4- dichlorophenoxyacetic acid) - BA (Benzyl adenine) - NAA (1-Naphthaleneacetic acid) - IAA (indole acetic acid) according to [3].

Table 1: Our Concentration from Hormones That Used

	Hormonal Transactions	Concentrations
1	BA + NAA	0.1 mg/l + 1 mg/l
2	BA + NAA	1 mg/l + 0.1 mg/l
3	BA + 2,4-D	0.3 mg/l + 0.75 mg/l
4	BA + 2,4-D	0.1 mg/l + 1 mg/l
5	BA + 2,4-D	2 mg/l + 1 mg/l
6	BA + 2,4-D	2 mg/l + 0.5 mg/l
7	BA + IAA	0.1 mg/l + 1 mg/l




Figure1: Induction of callus After 45 days from culture it was Observed Callus Induction with Good Grade by Using: (ba 0.1 mg/l+2.4-d1mg/l), (ba 0.3mg/l+ 2.4-d 0.75mg/l)



Figure 2: Light White to Yellowish Friable Callus Produced in Vitro Cultured Leaf Explants of Catharanthus Roseus at Ms Medium With Ba 0.3 Mg/L And 2,4-D 0.75 Mg/L Hormonal Combination.

Result and Discussion

These results show the variation and difference from the effect of different concentrations of hormones on tissue formation, as well as the effect of subcultures on the size and purity of the callus.




Figure3: Difference of Callus Formation When Used Different Concentration from

Table 2: Specific properties of *C. roseus*' TIAs and precursors for HPLC separation and detection, And the result from HPLC analysis was as follows: Ajmalicine (0.18 mg/g DW), Catharanthine (0.07 mg/g DW), Serpentine (0.37mg/g DW), Vindoline (0.11 mg/g DW).

Alkaloid	Fluorescence (max Ex/Em in nm)	Absorption UV (nm)
Tryptophan	250/330	198,231
Ajmalicine	250/350	217,265
Serpentine	310/380	272,367
Vindoline	201/261	201,239
Strictosidine	216/289	211,236
Catharanthine	129/181	160,197

Conclusion

The *Vinca rosea* plant contain indole alkaloids that have medical applications. High performance liquid chromatography was used with tandem mass spectrometry to determine Tryptophan, Ajmalicine, serpentine, vindoline, catharanthine, and Strictosidine. Overall, vinca alkaloids have the second most-used class of cancer drugs and will stay among the original cancer therapies. Different research and studies for new vinca alkaloid applications will be carried out in this regard. So, we conclude from this research that the different concentrations of growth hormones affect the differentiation and formation of callus where it was found that the use of hormonal BA and 2, 4-D is what gives the highest efficiency and vitality of the resulting callus [6-22].

References

- George EF, Hall MA, Klerk GJ De (2008) The Components of plant tissue culture media I: Macro- and micro-nutrients. Plant Propagation by Tissue Culture 3rd Edition 1: 65-113.
- Ghareeb Y, Soliman S, Ismail T, Hassan M (2016) Effect Of Growth Regulators And Ecotypes Of Egyptian *Vinca Rosa* (*Catharanthus Roseus*) On Callus Induction And Alkaloids Production. Zagazig Journal of Agricultural Research 43: 141-150.
- Swanberg A, Dai W (2008) Plant regeneration of periwinkle (*Catharanthus roseus*) via organogenesis. HortScience 43: 832-836.
- Ibrahim M, Ahmed MF, Kazim SM, Ghorri SS, Mehjabeen SS, et al. (2010) Antidiabetic activity of vinca rosea extracts in alloxan-induced diabetic rats. International Journal of

Endocrinology

- Sryea ATA, El-Ghit HMA (2012) Variation of Alkaloids Content of *Vinca Rosea* (*Catharanthus roseus* L) Under Different Meteorological Conditions of Five Governorates of Egypt 1: 60-68.
- Hoda E (2013) In Vitro Regeneration and Somaclonal Variation of *Catharanthus roseus* Don. Using Leaf and Internodal Explants. Alexandria Science Exchange Journal: An International Quarterly Journal of Science Agricultural Environments 34: 452-459.
- Meeting T, Fao J, Division I, Techniques N (2004) Low cost options for tissue culture technology in developing countries 26-30.
- Ataei-Azimi A, Delnavaz HB, Ebrahimzadeh H, Majd A (2008) High in vitro production of ant-canceric indole alkaloids from periwinkle (*Catharanthus roseus*) tissue culture. African Journal of Biotechnology 7: 2834-2839.
- Bakrudeen A, Subha Shanthi G, Gouthaman T, Kavitha M, Rao M (2011) In vitro micropropagation of catharanthus roseus-An anticancer medicinal plant. Acta Botanica Hungarica 53: 197-209.
- Faheem M, Singh S, Tanwer BS, Khan M (2011) In vitro Regeneration of multiplication shoots in *Catharanthus roseus* - An important medicinal plant. Palagia ResearchLibrary 2: 208-213.
- Kulkarni RN, Baskaran K, Jhang T (2016) Breeding medicinal plant, periwinkle [*Catharanthus roseus* (L) G. Don]: A review. Plant Genetic Resources: Characterisation and Utilisation 14: 283-302.
- Mishra JN, Verma NK (2017) A brief study on *Catharanthus Roseus* : A review. International Journal of Research in Pharmacy and Pharmaceutical Sciences 2: 20-23.
- Mishra MR, Akhtar N (2018) Enhancing alkaloid production from cell culture system of *Catharanthus roseus* with different carbon sources Enhancing alkaloid production from cell culture system of *Catharanthus roseus* with different carbon sources. 12-20.
- Sandhya M, Deepti L, Bhakti D, Ravindra M, Pranay S, et al. (2016) Effect of growth regulator combination on in-vitro regeneration of *Catharanthus roseus*. Int J Life Sci 6: 1-4.
- Alsanie S (2020) Molecular Studies on the Effect of some Auxins and Cytokinins on *Vinca Rosae* Grown by Plant Tissue Culture. Basic and Applied Sciences-Scientific Journal of King Faisal University.
- Asif M, Eudes F, Randhawa H, Amundsen E, Yanke J, et al. (2013) Cefotaxime prevents microbial contamination and improves microspore embryogenesis in wheat and triticale. Plant cell reports 32: 1637-1646.
- Javad S, Sumera Sarwar, Khajista Jabeen, Sumera Iqbal, Amina Tariq (2016) Enhanced extraction of an anticancer drug, Vinblastine, from *Catharanthus roseus*. Pure and Applied Biology 5: 608-614.
- Karimi M, F Raofie (2019) Micronization of vincristine extracted from *Catharanthus roseus* by expansion of supercritical fluid solution. The Journal of Supercritical Fluids 146: 172-179.
- Pietrosiuk A, et al (2007) *Catharanthus roseus*: micropropagation and in vitro techniques. Phytochemistry Reviews 6: 459-473.
- Salim V, Fang Yu, Joaquín Altarejos, Vincenzo De Luca (2013) Virus-induced gene silencing identifies *Catharanthus roseus* 7-deoxyloganic acid-7-hydroxylase, a step in iridoid and monoterpene indole alkaloid biosynthesis. Plant J 76: 754-765.

21. Siddiqui MJ, Z Ismail, NH Saidan et al (2011) Simultaneous determination of secondary metabolites from Vinca rosea plant extractives by reverse phase high performance liquid chromatography. Pharmacogn Mag 7: 92-96.
22. Zhang D, A Kanakkanthara (2020) Beyond the Paclitaxel and Vinca Alkaloids: Next Generation of Plant-Derived Microtubule-Targeting Agents with Potential Anticancer Activity. Cancers (Basel) 12: 1721.

Copyright: ©2022 Youssif Mahmoud Ahmed. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.