

## Research Article

## Open Access

## Evaluation of Serum Uric Acid Levels and Microalbuminuria in Newly Diagnosed Type 2 Diabetics with and without Hypertension

Dr Sujatha N Rao<sup>1</sup> and Kuldeep GB<sup>2</sup><sup>1</sup>Professor, Department of Biochemistry, AECS Maruthi College of Dental Sciences, Bengaluru, India<sup>2</sup>Chief Medical Administrator, Dept. of Medicine, Sri Krishna Sevashrama Hospital, Bngaluru, India**ABSTRACT**

**Objective:** This study aims to evaluate the levels of Serum uric acid (SUA) and microalbuminuria in Healthy, Newly Diagnosed (N.D.) type 2 Diabetics and in N.D. type 2 Diabetics with hypertension.

**Method:** Cross sectional study involving a total of 343 Individuals of Bengaluru, India from January 2018 to November 2019 were considered for the study. Venous blood and 24-hour urine sample were collected from 100 healthy, 170 N.D. type 2 Diabetics, 73 N.D. type 2 Diabetics with hypertension and analyzed for Glycated hemoglobin (HbA1c) SUA and microalbuminuria along with their demographic details.

**Results:** There was significant increase in SUA levels and microalbuminuria in N.D. Type 2 Diabetics with hypertension compared to healthy and N.D. type 2 Diabetics.

**Conclusion:** Hypertensive Diabetics should be monitored periodically for SUA and microalbuminuria to prevent them from developing renal complication.

**\*Corresponding author**

Dr Sujatha N Rao, Professor, Department of Biochemistry, AECS Maruthi College of Dental Sciences & Research Center off Bannerghatta Road, Bengaluru, India, 560076; Email: sujatharao.8@gmail.com

**Received:** October 10, 2022; **Accepted:** October 17, 2022; **Published:** October 25, 2022

**Keywords:** Glycated Hemoglobin, HbA1c, Hypertension, Newly Diagnosed (N.D) type 2 Diabetics, Serum uric acid, Microalbuminuria.

**Introduction**

Type 2 Diabetes Mellitus (DM) characterized by Hyperglycemia due to inefficient insulin is one of the most important public health concern worldwide [1-4]. It is predicted that the countries with largest no. of people with diabetes mellitus will be from India, China, and USA in the year 2025 [5, 6]. Hyperglycemia can be determined by Glycated Hemoglobin (HbA1c) and Fasting Glucose (FBS). HbA1c reflects the blood glucose level over the previous 3 months [7]. Uric Acid is a product of purine catabolism. It is a powerful plasma soluble antioxidant within its normal range. 2-7 mg/dl in Men and 2-6 mg /dl in women [8]. Although several Investigators have reported SUA as biological marker for type 2 diabetes, their study results were contradictory. A few Researchers through their studies on type 2 diabetic population in USA, Sweden and Netherlands have reported elevated SUA [9-12]. Many Investigators have reported low serum uric acid levels within the Chinese type 2 Diabetic population [13-16]. While a few others have reported elevated serum uric acid levels [17, 18]. In addition, within the Japanese type 2 Diabetic population, few investigators have reported lowered SUA while a few other investigators have

reported elevated SUA [19, 20]. Microalbuminuria is an important predictor of progression to overt proteinuria which results in decreased GFR that may progress into End Stage Renal Disease (ESRD) gradually [21, 22]. Due to these reasons, our study aims to evaluate SUA levels and microalbuminuria in N.D. Type 2 Diabetic Individuals with and without hypertension and to compare these parameters with those of Healthy population.

**Materials and Methods**

This was a cross sectional study conducted at the Shree Krishna Sevashrama Hospital, Bengaluru, India between January 2018 to November 2019. A total of 343 Adult Individuals were included in this study, in the age group 38 to 62 years of both the gender. After obtaining the Institutional Ethical committee approval, informed consent was taken from all the individuals participated in the study. Individuals were categorized into 3 groups, based on their Hb A1c levels as per W.H.O. criteria [23, 24].

**Inclusion Criteria**

Group I included 100 healthy individuals with HbA1c equal to or below 5.6

Group II included 170 N.D. type 2 Diabetics with HbA1c level 6.5 or above

Group III included 73 N.D. type 2 Diabetics with hypertension

with HbA1c level 6.5 or above and with blood pressure 140/90 or above.

Exclusion Criteria for all the above 4 groups included pregnant and lactating women, individuals taking medications.

### Collection of Sample

After an overnight fast, blood samples were obtained for biochemical analysis. Sample collection involved venous whole blood sample. Portion of it was transferred to EDTA tubes for the analysis of HbA1c and FBS and the other portion was allowed to clot. Serum was separated and used for the analysis of SUA. 24-hour urine sample was also collected for detection and estimation of microalbuminuria.

### Analysis

Analysis was performed using Olympus AV Auto analyser, using Diasys reagents manufactured by Diasys Diagnostic system, GmbH, Holzheim, Germany.

HbA1c was measured using particle enhanced Immuno turbidimetric method and the desirable range for healthy population was  $\leq 5.6$ , for prediabetes 5.7 – 6.4, for diabetes,  $\geq 6.5$  [25]. SUA was analysed by enzymatic method and the desirable range for SUA was 2-7 mg/dl in men and 2-6 mg /dl in women [26]. Result of all the above parameters were expressed as mean with standard deviation.

### Statistical Analysis

Data were analyzed using SPSS statistical version 24. One way Analysis of Variance (ANOVA) was performed to compare the means of the three groups. Statistical difference between any 2 groups among the total 3 groups was measured by 't' test. Results of all the tests with  $p < 0.05$  were considered statistically significant and highly significant at  $P < 0.01$ .

### Results

**Table No. I – Serum Uric Acid Levels in Three Different Populations.** Significant at  $P < 0.05$ .

Highly significant at  $P < 0.01$

Serum uric acid mg/dl	N	Mean	Std Deviation	F	Sig.
Healthy	100	4.212	0.589	249.510	0.000
N.D. Type 2 Diabetics	170	4.119	0.621		
N.D. Type 2 Diabetics with Hypertension	73	7.730	0.876		

**Table No.2- Microalbuminuria in Three Different Populations.** Significant at  $p < 0.05$ . Highly significant at  $P < 0.01$

Microalbuminuria mg/24 hours	N	Mean	Std. Deviation	F value	Sig. P
Healthy	100	14.20	1.99	363.804	0.000
N.D. type 2Diabetics	170	16.96	2.03		
N.D. type 2 Diabetics with hypertension	73	45.61	38.35		

**Table No. 3 Comparison of SUA Levels between the above 2 Groups.** Significant at  $P < 0.05$ . Highly significant at  $P < 0.01$ .

Group	N	Mean	Std Deviation	t	Sig.
Serum uric acid mg/dl					
N.D. type 2 Diabetics with Hypertension	73	7.730	0.876	17.856	0.000
N.D. type 2 Diabetics	170	4.319	0.621		

**Table No. 4 Comparison of SUA Levels between the above 2 Groups.** Significant at  $P < 0.05$ . Highly significant at  $P < 0.01$ .

Group	N	Mean	Std Deviation	t	Sig.
Serum uric acid					
Diabetics with Hypertension	73	7.730	0.876	20.741	0.000
mg/dl	100	4.212	0.589		

**Table No. 5 Comparison of SUA levels between the 2 Groups.** Significant at  $P < 0.05$ . Highly significant at  $P < 0.01$

Group	N	Mean	Std Deviation	t	Sig.
Uric acid					
Healthy	100	4.212	0.589	0.38	0.790
mg/dl	170	4.319	0.621		

**Table No.6 Comparison of Microalbuminuria between the 2 Groups.** Significant at  $P < 0.05$ . Highly significant at  $P < 0.01$

	Group	N	Mean	Std Deviation	t	Sig.
Microalbuminuria	Healthy	100	14.20	1.99	0.55	0.727
mg/24 hours	N.D. Diabetics	170	16.96	2.03		

**Table No.7 Comparison of Microalbuminuria between the 2 Groups.** Significant at  $P < 0.05$ . Highly significant at  $P < 0.01$ .

	Group	N	Mean	Std Deviation	t	Sig.
Microalbuminuria	N.D. Diabetics	170	16.96	2.03	56.240	0.000
mg/24 hours	N.D. Diabetics with hypertension	73	45.61	38.35		

**Table No. 8 Comparison of Microalbuminuria between the 2 Groups.** Significant at  $P < 0.05$ . Highly significant at  $P < 0.01$

	Group	N	Mean	Std Deviation	t	Sig.
Microalbuminuria	Healthy	100	14.20	1.99	56.937	0.000
mg/dl	N.D. Diabetics with hypertension	73	45.61	38.35		

Table 1 displays the level of SUA in 3 different populations. Among these, only in N.D. Diabetics with hypertension group, SUA is above the normal range ( $7.730 \pm 0.876$ ). SUA levels are within the normal range in Healthy group ( $4.212 \pm 0.589$ ) and in N.D. Type 2 Diabetics group ( $4.319 \pm 0.621$ ). Significant difference ( $P=0.000$ ) is noted in SUA levels between N.D. Diabetics group and N.D. Diabetics with Hypertension groups (Table 3). Significant difference ( $P=0.000$ ) is noted in SUA levels between Healthy group and N.D. Diabetics with Hypertension

groups (Table No. 4). No significant difference ( $P=0.790$ ) is noted in SUA levels between Healthy group and N.D. Diabetics group (Table 5)

Table 2 displays the amount of microalbuminuria in 3 different populations.) Increased Microalbuminuria was noted only in N.D. type 2 Diabetics with hypertension group. No significant difference ( $P=0.727$ ) is noted in microalbuminuria between Healthy group and N.D. Diabetics group (Table 6). Significant difference ( $P=0.000$ ) is noted in microalbuminuria between N.D. Diabetics group and N.D. Diabetics with Hypertension groups (Table 7). Significant difference in ( $P=0.000$ ) is noted in Microalbuminuria levels between Healthy group and N.D. Diabetics with Hypertension groups (Table No. 8).

## Discussion

We have observed elevated SUA levels above the normal range only in N.D. type 2 Diabetics with hypertension population and not in N.D. type 2 diabetics population. Reason may be that our diabetic population is a newly diagnosed one. Elevated SUA in N.D. type 2 Diabetics with hypertension may be due to the defect in the urate transporter, which leads to enhanced renal reabsorption of uric acid resulting in elevated SUA levels. In addition, we have noted significantly increased microalbuminuria in N.D. type 2 Diabetics with hypertension compared to the other 2 Groups. There was neither significant increase in its amount in the other 2 groups nor significant difference in the amount of microalbuminuria between the same 2 groups. Increased microalbuminuria in N.D. type 2 Diabetics with hypertension group may be an indicator of the condition progressing into Diabetic Nephropathy due to increased vascular permeability and endothelial dysfunction [21, 22].

## Conclusion

Elevated SUA and increased microalbuminuria were observed in N.D. type 2 diabetics with hypertension population. Hence, periodic monitoring of these two parameters while monitoring the case of hypertensive diabetics may prevent their condition from developing renal complication. In addition, early detection and timely action reduces the economic burden associated with Diabetes in developing countries.

## Acknowledgements

The authors thank Sri. Krishna Sevashrama Hospital, Bengaluru for granting an opportunity to conduct this study. A special thanks to the Nursing Superintendent Mrs. Judith Fernandes for her help in recording the data.

## Disclosure Statement

The authors have no conflict of interest to declare.

## References

1. Bastaki S (2005) Diabetes mellitus and its treatment. *International Journal of Diabetes and Metabolism* 13: 111-134.
2. Liamis G, Liberopoulos E, Barkas F, Elisaf M (2014) Diabetes mellitus and electrolyte disorders. *World Journal of Clinical Cases* 2: 488.
3. Paul Z Zimmet (2017) Diabetes and its Drivers: the largest epidemic in human history. *Clin.diabet.endocrinol* 3: 1.
4. Tandon N, Anjana RM, Mohan V, Kaur T, Afshin A (2018) The increasing burden of Diabetes and variations among the states of India: the Global Burden of disease study 1990-2016. *Lancet Global Health* 6: 1352-1362.
5. Lei Chen, Dianna J. Magliano, Paul Z. Zimmet (2012) The worldwide epidemiology of type 2 diabetes mellitus-present and future perspectives. *Nature Reviews Endocrinology* 8: 228-236.
6. LGSR Mehta, Kashyap AS, Das S (2009) Diabetes Mellitus in India; the Modern Scourge. *MJAFI* 65: 50-54.
7. Alam R, Verma MK, Verma P (2015) Glycated Hemoglobin as a dual Biomarker in Type 2 Diabetes Mellitus predicting glycemic control and dyslipidemia risk. *TC* 189: 164-189.
8. Maiuolo J,oppedisano F, Gratteri S, Muscali C, Mollace V (2016) Regulation of Uric acid Metabolism and Excretion. *International Journal of Cardiology* 213: 8-14.
9. Bhandaru P, Shankar A (2011) Association between serum uric acid levels and Diabetes mellitus. *International Journal of Endocrinology* 7604715: 1-7.
10. Ohlson LO, Larsson B, Bjorntorp P, Eriksson H, Svärdsudd K, et al. (1988) Risk factors for type 2 Diabetes Mellitus: Thirteen and one half years of follow up of the participants in a study of Swedish Men born in 1913. *Diabetologica* 11: 798-805.
11. Dehgan A, Hock M Van, Sijbrands EJG, Hofman A, Wetteman JCM (2008) High Serum Uric acid level as a Novel Risk factor for type 2 Diabetes. *Cardiovasc. Metab. Risk* 31: 361-362.
12. Krammer CKD, Von Muhlen, Jassal SK, E-Barrett-connor (2009) Serum uric acid Levels improve predictions of incident type 2 Diabetes in Individuals with Impaired fasting glucose: The Rencho Bernado study. *Diabetic care* 32: 1272-1273.
13. Nan H, Dong Y, Gao W, Tuo Milehto J, Qiao Q (2007) Diabetes associated with low serum uric acid level in general Chinese population. *Diabetes Res.Chin. Pract* 76: 68-74.
14. Wei F, Chang B, Yang X, Wang Y, Chen L, et al. (2016) Serum uric acid levels were Dynamically coupled with Hemoglobin A/C in the development of Type 2 Diabetes. *Sci. Rep* 22: 28549.
15. Hairong Nan, Zengcheng pan, Shaojewang, Weighogao, Lei Zhang, et al. (2009) Serum uric acid, Plasma Glucose and Diabetes. *Diabetes Vascular Diseases Research* 7: 40-46.
16. Yue B, Tan JB, Ning F, Sun JP, Zhang KY, et al. (2015) Association Between Serum uric acid and prevalence of Type 2 Diabetes Diagnosed using Hb A/C criteria among Chinese adults in Ringdo, china. *Biomed Environ-Su* 12: 884-893.
17. Yuliang Cui, Hemei Bu, Xianghua Ma, Sha Zhao, Xiaona Li, et al. (2016) The Relationship between SUA and Hb A1C is dependent upon Hyper insulinemia in patients with newly diagnosed type 2 Diabetes mellitus. *Diabetes Res* 2016: 1-7.
18. WYN YT, chan CSY, Lui SS (1999) Hyperurecemia in type 2 Diabetes. *Diabetes Nutr. Metab* 12: 286-291.
19. E Oda, R Kawai, V Sukumaran, K Watanabe (2009) Uric acid is positively associated with metabolic syndrome but negatively associated with Diabetes in Japanese Men. *Internal Medicine* 48: 1785-1791.
20. Nakanishi N, Okamoto M, Yoshida H Matsuo Y, Suzuki K, Tataru K (2003) Serum uric acid and risk development of Hypertension and impaired fasting glucose or type II Diabetes in Japanese male office workers. *Eur. J. Epidemiology* 18: 523-530.
21. Fauci AS, Braunwald E, Kasper DL, Hauser SL, Longo DL, et al. (2008) Principles of Internal Medicine, Harrison's 17th edition. Mc Graw-Hill. [https://muhammad1988adeel.files.wordpress.com/2011/10/harrison\\_\\_s\\_-\\_manual\\_of\\_medicine\\_-\\_17th\\_edition.pdf](https://muhammad1988adeel.files.wordpress.com/2011/10/harrison__s_-_manual_of_medicine_-_17th_edition.pdf).
22. Basi S, Fesler P, Mimran A, Lewis JB (2008) Microalbuminuria in type 2 diabetes and hypertension: a marker, treatment target, or innocent bystander?. *Diabetes Care* 31: S194-S201
23. American Diabetes Association (2018) Classification and

- diagnosis of diabetes: standards of medical care in diabetes. Diabetes Care 41: S13-27.
24. The International Expert committee (2009) International Expert Committee report on the role of A1c assay in the diagnosis of diabetes. Diabetes care 32: 1327-1334.
25. Nordin G (2007) Analysis of HbA1C by particle Enhanced Immunoturbidimetric method.
26. Fossati P, Prencipe L, Berti G (1980) Use of 3,5-dichloro-2-hydroxybenzenesulfonic acid/4-aminophenazone chromogenic system in direct enzymic assay of uric acid in serum and urine. Clin Chem 26: 227-231.