

Invasive Potential of Nile Tilapia, *Oreochromis Niloticus* (Linnaeus, 1758) From the Tributary of the Ganga River, Central India

Priyanka Mayank¹, Neeti Mishra¹ and Amitabh Chandra Dwivedi^{2*}

¹Kalash Research and Welfare Society, Prayagraj-211002, Uttar Pradesh, India

²Department of Zoology, Nehru Gram Bharati (Deemed to be University), Prayagraj, Uttar Pradesh, India

ABSTRACT

The natural fishery or wild stock of fishes especially fresh water in India is significantly declining due to manmade stressors. The invasion of fishes and size composition are an essential component of stream, river and other large water bodies ecosystem and represents an evident of structure, function, depth and health of stream/river. During study period 683 specimens of *Oreochromis niloticus* were collected from February 2019 to January 2020 in Sirsa fish landing centre from the lower stretch of the Tons river at Prayagraj, Uttar Pradesh, India. Size composition of fishes was varied from 82 to 463 mm (total length). The maximum total length of *O. niloticus* in the catches indicated that its increasing colonisation success in the river Tons. Size composition of fishes was also indicated that the stock of *O. niloticus* in the Tons river was in healthy condition. The 231-260 mm size group was most dominated (15.67%) compared to 261-290 mm (12.44%) and 291-320 mm (11.71%) in the total exploited stock. The middle size group was maximum exploited with 51.24% from the Tons river at Prayagraj. Higher size group contributed sizeable proportion with 25.04% in the exploited stock. The exploitation pattern was unsystematic in higher size group due to contribution of higher proportion. Current exploitation pattern is unsystematic but *O. niloticus* is powerfully invaded from the lower stretch of the Tons river at Prayagraj.

*Corresponding author

Amitabh Chandra Dwivedi, Department of Zoology, Nehru Gram Bharati (Deemed to be University), Prayagraj, Uttar Pradesh, India. Tel: +919450211911; E-mail: saajjan@rediffmail.com

Received: June 14, 2021; Accepted: June 22, 2021; Published: June 30, 2021

Keywords: *Oreochromis Niloticus*, Size Composition, Invasive Potential, Exploitation, Tons River

Introduction

Fishes are significant to providing a high-value protein source with omega-3 fatty acid and important micronutrients for much of the globe [1]. The human activities have powerfully altered size, age, sex ratio, fecundity, feeding nature, exploitation and biodiversity of fishes especially from the freshwater ecosystems over the world [2-9]. In general, *Oreochromis niloticus* (Nile tilapia) is eminent for their plasticity in survival, feeding nature, fast growth, tolerance, ease of breeding and size-at first maturity that not only compose them the ability to invasion in any fresh water ecosystem and become established in non-native ecosystem or introduced ecosystem (example habitats, environments) but also allow them an excellent aquaculture species (Example due to its ability to tolerate a wide range of environmental conditions and ability to feed at different trophic levels) with other carps [10-16]. It is naturally distributed in the Nile River as well as most parts of African rivers, reservoirs and lakes while it is exotic fish species for India [17-20]. It is a great economical importance species and significant role in the tropical and sub-tropical aquatic ecosystems [21-23]. Many researchers have reported that the invasive cichlids (Example *Oreochromis niloticus*) reducing growth and reproduction especially fecundity and survival of spawn of native species [24-28].

In 20th century, *O. niloticus* has been intentionally dispersed worldwide, in particular for aquaculture, restocking programs and higher production [17,18,29-33]. As production from capture fisheries (Example riverine sector) decreasing day by day but invasion of *Oreochromis niloticus* (Nile tilapia) and *Cyprinus carpio* (Common carp) it increased from the Ganga river system [34-36]. It is commercially and economically exploited from the Ganga river with 14.56% in 2015-2016 and Yamuna river with 24.36% in 2011-2012 at Allahabad (now Prayagraj) [19,37]. Therefore, knowledge of size composition and exploitation pattern of *O. niloticus* is essential to better understand its stock status, impact and to design optimal management strategies. Presently, the Tons river (lower stretch) fishery is very healthy in respect of landing, and size composition and exploitation pattern investigations are timely as the data are needed to improved management of *O. niloticus* stock and impact in the fishery of Indian major carp (Catla catla, Labeo rohita, Cirrhinus mrigala) and large size carp and catfishes (Labeo calbasu, Sperata aor, S. seenghala, Wallago attu).

Material and methods

Climate and Characteristics of the River

The climate of this region (Tons river basin) is marked by mild cold during winter and intensive heat during summer. The monsoon season is July to September month. Sometimes winter rainfall is also recorded. The Tons river is essentially a hilly stream arising in the Kaimur hills of the Vindhyan range, Madhya Pradesh, India. The

Tons river drain the Bundelkhand geographic region of central India. Bundelkhand lies between the Indo-Gangetic Plain to the north and the Vindhya Range to the south. It is a tributary of the Ganga, which forms confluence at Sirsa near Meja in the Prayagraj district. Tons river lies between latitude 24° 0' to 25° 16' 54" North and longitude 80° 26' 45" to 82° 04' 57" East. Its banks are lined by deep ravines and the bed is rocky. Agriculture and human settlements were the major land use category in its catchment.

During study period 683 specimens of *Oreochromis niloticus* were collected from February 2019 to January 2020 in Sirsa fish landing centre from the lower stretch of the Tons river at Prayagraj, Uttar Pradesh, India. Size composition (total length) varied from 82 to 483 mm. Drag net, cast net, gill net and hook and line were used by fishers/fishermen to catch the fishes in the Tons river. A total of 683 fish samples (male and female) were collected and analyzed. The total length (mm) from the tip of snout to the end of largest caudal fin rays was measured by measuring scale. The obtained data from the river was classified into a series of size groups of 30 mm intervals. The number of samples calculated according to size group then converted into percentage.

Result and discussion

The size composition of *O. niloticus* was varied from 82 mm to 463 mm (total length) from the lower stretch of the Tons river at Prayagraj, Uttar Pradesh, India. The maximum fish exploitation was recorded in 231 to 260 mm size group in case of male, female and pooled samples. This size group of fishes is more attracted to the fishers (Example due to market value, cost of fishing) and consumers also (Example taste, structure of bone in fish body) both. The maximum total length (463 mm) of *O. niloticus* in the catch indicated that its increasing colonisation success from the river Tons at Prayagraj, Uttar Pradesh, India.

Pooled samples

The maximum fish exploitation was recorded in 231 to 260 mm size group with 15.67% while minimum exploitation was observed with 1.17% in 441-460 mm size group. Fish exploitation is an economic activity governed by social needs and pressures. The middle size group was maximum exploited compared to lower and higher size groups from the lower stretch of the Tons river at Prayagraj. The experienced mature female fish stock was healthy in the river in monsoon season but very high fishing pressure we observed in this season. On the basis of data, it is observed that middle size group was maximum exploited with 51.24% at Prayagraj. Lower size group was exploited with 23.78%. Higher size group shared sizeable proportion with 25.04% in exploited population (Figure 1). The results also indicated that the exploitation was unsystematic in higher size group.




Figure 1: Exploitation structure (according to size group) of *Oreochromis niloticus* from the Tons river at Prayagraj, Uttar Pradesh

Lower size groups 81-110 mm, 111-140 mm 141-170 mm and 171-200 mm were contributed in the exploited stock with 2.34%, 5.41%, 7.17% and 8.78%, respectively (Table 1). Middle size groups 201-230 mm, 231-260 mm 261-290 mm and 291-320 mm were shared in the exploitation 11.42%, 15.67%, 12.44% and 3.28%, respectively. Higher size groups 321-350 mm, 351-580 mm, 381-410 mm, 411-440 and 441-470 mm were contributed in the exploitation with 10.54%, 7.03%, 4.54%, 1.76% and 1.17%, respectively (Table 1, Figure 2).




Figure 2: Size composition and exploitation structure of *Oreochromis niloticus* from the Tons river at Prayagraj, Uttar Pradesh (Pooled samples)

Male samples

In case of male samples, the maximum exploitation was recorded in 231 to 260 mm size group with 15.18% but minimum exploitation was observed with 1.49% in 441-460 mm size group (Table 1). Size group 261-290 mm was the second most exploited stock (Fig. 3). Size group 81-110 mm, 111-140 mm 141-170 mm and 171-200 mm, 201-230 mm, 291-320 mm, 321-350 mm, 321-350 mm, 351-380 mm, 381-410 mm and 411-440 mm were contributed in the exploitation with 2.68%, 5.95%, 7.73%, 9.23%, 11.01%, 11.61%, 10.12%, 6.84%, 4.17% and 2.08%, respectively (Figure 3).




Figure 3: Size composition and exploitation structure of *Oreochromis niloticus* from the Tons river at Prayagraj, Uttar Pradesh (Male samples)

Female samples

In case of female samples, the maximum exploitation was recorded in 231 to 260 mm size group with 15.67% while minimum exploitation was observed with 1.17% in 441-460 mm size group (Table 1). Size group 261-290 mm was the second most exploited stock (Figure 4). Size group 81-110 mm, 111-140 mm 141-170 mm and 171-200 mm, 201-230 mm, 291-320 mm, 321-350 mm and 321-350 mm were shared in the exploitation with 2.68%, 5.95%, 7.73%, 9.23%, 11.01%, 11.61% and 10.12%, respectively (Fig. 4). 351-380 mm, 381-410 mm and 411-440 mm size group was contributed minute proportion with 6.84%, 4.17% and 2.08%,

respectively. Reported 41.3 to 400.0 mm (total length) size of fishes, *O. niloticus* in coastal Mississippi, USA [18]. Estimated that the total length 13.5 to 48.0 cm of *Oreochromis niloticus* from Lake Nasser, Egypt [38,39]. Rreported that the size composition of *O. niloticus* varied from 10.4 to 44.5 cm from the Yamuna river at Allahabad (now Prayagraj), Uttar Pradesh. She also stated that the aquatic environment of the Yamuna river at Allahabad was most favorable for *O. niloticus* due to fast growth increment, food supply and multiple breeding. Reported that the frequency of fishes of age group I are most dominant in the catch and constitute about 33.49%, 36.42% and 33.74% for *O. niloticus* males, females and sexes combined, respectively in El-Bahr El-Faraouny Canal, Al-Minufiya Province, Egypt [40].




Figure 4: Size composition and exploitation structure of *Oreochromis niloticus* from the Tons river at Prayagraj, Uttar Pradesh (Female samples)

Fishing pressure changes the size composition, total length, biodiversity, density of fishes, growth rate, age composition, sex ratio, income of fishers/fishermen and maturation of fishes [41-47]. The size composition (Example maximum and minimum total length) of fishes is altering season to season and year to year due to large size fishes more active in breeding season [48-51]. The over exploitation and non-targeted fishing is the biggest problem of riverine fishery [52-53]. The fishing pressure, mesh size, size of nets and fishing technique (example degree) are responsible for increasing or decreasing of total length of fishes (example maximum total length) and recruitment in the lotic ecosystems [47, 54-57]. Non-native fishes are also changed selectivity of gear due to nature, dwelling behaviour and ecological condition [58-59]. The growth of fishes is slightly checked by heavy metals accumulation in the body of fishes [60-62].

Conclusion

It may be concluded that the research provides an important baseline study of this fish. Size composition of *O. niloticus* indicated that the stock from the Tons river was in healthy condition and exploitation was unsystematic. Overall, present condition of *O. niloticus* was recorded due to sustainable exploitation and multiple breeding of this species in the Tons river. Mostly large size and perennial rivers has large size of fishes in water bodies and maintain food supply, space, reproductive needs stated that the when sustainably harvested or farmed, inland fish can be considered part of the green food movement for more environmentally friendly sourcing of food [41, 53, 63-65].

Table 1: Size composition and exploitation of Nile tilapia, *Oreochromis niloticus* from the Tons river at Prayagraj, Uttar Pradesh, India

SN	Size groups (mm)	Male	%	Female	%	Pooled	%
1	81-110	09	2.68	07	2.02	16	2.34
2	111-140	20	5.95	17	4.90	37	5.41
3	141-170	26	7.73	23	6.63	49	7.17
4	171-200	31	9.23	29	8.36	60	8.78
5	201-230	37	11.01	41	11.81	78	11.42
6	231-260	51	15.18	56	16.14	107	15.67
7	261-290	40	11.90	45	12.97	85	12.44
8	291-320	39	11.61	41	11.81	80	11.71
9	321-350	34	10.12	38	10.95	72	10.54
10	351-380	23	6.84	25	7.20	48	7.03
11	381-410	14	4.17	17	4.90	31	4.54
12	411-440	07	2.08	05	1.44	12	1.76
13	441-470	05	1.49	03	0.86	08	1.17
	Total	336	49.19	347	50.80	683	

References

- Mishra N, Dwivedi AC (2020) Environmental drivers supports to distribution, composition and biology of *Cyprinus carpio* (Linnaeus, 1758) in respect of time scale: A review. Journal of the Kalash Science 8: 91-102.
- Peterson MS, Slack WT, Waggy GL, Finley J, Woodley CM, et al. (2006) Foraging in non-native environments: comparison of Nile Tilapia and three co-occurring native centrarchids in invaded coastal Mississippi watersheds. Environmental Biology of Fishes 76: 283-301.
- Arlinghaus R, Lorenzen K, Johnson BM, Cooke SJ, Cowx IG (2015) Management of freshwater fisheries: addressing habitat, people and fishes. In: Freshwater fisheries ecology. edited by J. F. Craig. John Wiley & sons Ltd. 1st Ed 557- 579.
- Lynch AJ, Cooke SJ, Deines AM, Bower SD, Bunnell DB, et al. (2016) The social, economic and environmental importance of inland fish and fisheries. Environ. Rev., doi: 10.1139/er-2015-0064.
- Lacy SN, Meza FJ, Marquet PA (2017) Can environmental impact assessments alone conserve freshwater fish biota? Review of the Chilean experience. Environmental Impact Assessment Review 63: 87-94.
- Dwivedi AC, Mishra AS, Mayank P, Tiwari A (2016) Persistence and structure of the fish assemblage from the

- Ganga river (Kanpur to Varanasi section), India. Journal of Geography and Natural Disasters 6: 159.
7. Dwivedi AC, Mishra AS, Mayank P, Tripathi S, Tiwari A (2019) Resource use competence and invader potential of Cyprinus carpio from the Paisuni river at Bundelkhand region, India. Journal of Nehru Gram Bharati University 8: 20-29.
 8. Ripple WJ, Wolf C, Newsome TM, Bernard P, Moomaw (2020) World scientists warning of a climate emergency. BioScience 70: 8-12
 9. Mishra N, Dwivedi AC, Mayank P (2021) Invasion potential, impact and population structure of non-native fish species, Cyprinus carpio (Linnaeus, 1758) from the tributary of the Ganga River, Central India. Aquaculture and Fisheries Studies 3: 1-4.
 10. Zambrano L, Martinez-Meyer E, Menezes N, Peterson AT (2006) Invasive potential of common carp (Cyprinus carpio) and Nile tilapia (*Oreochromis niloticus*) in American freshwater systems. Canadian J Fisheries Aquatic Sci 63: 1903-1910.
 11. Ishikawa T, Shimose T, Tachihara K (2013) Life history of an invasive and unexploited population of Nile tilapia (*Oreochromis niloticus*) and geographical variation across its native and non-native ranges. Environ. Biol. Fish 96: 603- 616.
 12. Alam A, Chadha NK, Joshi KD, Chakraborty SK, Sawant PB, et al. (2015) Maturation profile and fecundity of the exotic *Oreochromis niloticus* in the river Yamuna, India. Journal of Environmental Biology 36: 927-93.
 13. Dwivedi AC, Mayank P (2018) Suitability of ecosystem determination through biology and marketing of exotic fish species, *Oreochromis niloticus* (Linnaeus, 1757) from the Ganga River, India. Journal of Aquatic Research and Marine Sciences 1: 69-75
 14. Tiwari A, Dwivedi AC, Mayank P (2016) Time scale changes in the water quality of the Ganga River, India and estimation of suitability for exotic and hardy fishes. Hydrology Current Research 7: 254.
 15. Abarike ED, Ampofo-Yeboah A (2016) Reproductive potential of Nile tilapia (*Oreochromis niloticus* Linnaeus, 1757) in the Gologing reservoir in Ghana. International Journal of Fisheries and Aquatic Studies 4: 279-283.
 16. Tripathi S, Gopesh A, Dwivedi AC (2017) Fish and fisheries in the Ganga river: current assessment of the fish community, threats and restoration. J Exp Zoology 20: 907-912.
 17. Trewavas E (1983) Tilapiine fishes of the genera Sarotherodon, Oreochromis and Danakilia. London: British Museum Natural History.
 18. Grammer GL, Slack WT, Peterson MS, Dugo MA (2012) Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758) establishment in temperate Mississippi, USA: multi-year survival confirmed by otolith ages. Aquatic Invasions 7: 367-376.
 19. Tripathi S, Gopesh A, Dwivedi AC (2017) Framework and sustainable audit for the assessing of the Ganga river ecosystem health at Allahabad, India. Asian Journal of Environmental Science, 12: 37-42.
 20. Dwivedi AC, Mayank P, Tiwari A (2016) The River as transformed by human activities: the rise of the invader potential of Cyprinus carpio and *Oreochromis niloticus* from the Yamuna River, India. Journal of Earth Science & Climatic Change 7: 361.
 21. Bucur C, Costache M, Daniel O, Nino M (2012) Studies and observations on the spawning of *Oreochromis niloticus* species reared at SCDP Nucet-Dambovita. Animal Science and Biotechnologies 45: 1-6.
 22. Tsungai AZ, Robertson MP, Booth AJ, Chimimba CT (2013) A qualitative ecological risk assessment of the invasive Nile tilapia, *Oreochromis niloticus* in a sub-tropical African river system (Limpopo River, South Africa). Aquatic Conservation: Marine and Freshwater Ecosystems 23: 51-64.
 23. Mayank P, Dwivedi AC (2017) Resource use efficiency and invasive potential of non-native fish species, *Oreochromis niloticus* from the Paisuni River, India. Poultry Fisheries & Wildlife Sciences 5.
 24. Lorenz OT, O'Connell MT (2008) Growth of non-native Rio Grande cichlids (*Herichthys cyanoguttatus*) at different salinities and in the presence of native bluegill *Lepomis macrochirus*. Journal of Freshwater Ecology 23: 537-544.
 25. Doupé, RG, Schaffer J, Knott MJ, Burrows DW (2009) How might an exotic fish disrupt spawning success in a sympatric native species? Marine and Freshwater Research 60: 379-383.
 26. Gomez-Marquez JL, Pena-Mendoza B, Salgadougarte IH, Arrondo-Figueroa JL (2008) Age and growth of the tilapia, *Oreochromis niloticus* (Perciformes: Cichlidae) from a tropical shallow lake in Mexico. Rev. Biol. Trop 56: 875-884.
 27. Lorenz OT, O'Connell MT, Schofield PJ (2011) Aggressive interactions between the invasive Rio Grande cichlid (*Herichthys cyanoguttatus*) and native bluegill (*Lepomis macrochirus*), with notes on redspotted sunfish (*Lepomis miniatus*). Journal of Ethology 29: 39-46.
 28. Mayank P, Dwivedi AC (2015) Role of exotic carp, Cyprinus carpio and *Oreochromis niloticus* from the lower stretch of the Yamuna river. In: Advances in Biosciences and Technology Edited by K. B. Pandeya, A. S. Mishra, R. P. Ojha and A. K. Singh published by NGB (DU), Allahabad 93-97.
 29. Canonico GC, Arthington A, McCrary JK, Thieme ML (2005) The effects of introduced tilapias on native biodiversity. Aquat Conserv Mar Freshwat Ecosyst 15: 463-83.
 30. Schofield PJ, Slack WT, Peterson MS, Gregoire DR (2007) Assessment and control of an invasive aquaculture species: an update on Nile tilapia (*Oreochromis niloticus*) in coastal Mississippi after Hurricane Katrina. Southeastern Fishes Council Proceedings 49: 9-15.
 31. Schofield PJ, Peterson MS, Lowe MR, Brown-Peterson N, Slack WT (2011) Survival, growth and reproduction of nonindigenous Nile tilapia (*Oreochromis niloticus* (Linnaeus, 1758)). I. Physiological capabilities to various temperatures and salinities. Marine and Freshwater Research 62: 439-449.
 32. Gozlan RE, Britton JR, Cowx I, Copp GH (2010). Current knowledge on non-native freshwater fish introductions. Journal of Fish Biology 76: 751-786.
 33. Dwivedi AC, Mayank P, Tripathi S, Tiwari A (2017) Biodiversity: the non-natives species versus the natives species and ecosystem functioning. Journal of Biodiversity, Bioprospecting and Development 4.
 34. Jha DN, Joshi KD, Dwivedi AC, Mayank P, Kumar M, et al. (2015) Assessment of fish production potential of Chitrakoot district, Uttar Pradesh. Journal of the Kalash Science 3: 7-10.
 35. Alam A, Chadha NK, Chakraborty SK, Joshi KD, Das SCS, et al. (2019) Studies on the growth and mortality of invasive *Oreochromis niloticus* (Linnaeus, 1758) in sub-tropical river Yamuna, part of Gangetic River system, India. Aquatic Ecosystem Health & Management, DOI: 10.1080/14634988.2019.1690926.
 36. Dwivedi AC, Mishra N (2021) Age structure of non-native fish species, Cyprinus carpio (Linnaeus, 1758) from the tributary of the Ganga river, India. Journal of Aquaculture & Marine Biology 10: 76-79.
 37. Mayank P, Dwivedi AC (2015) Biology of Cirrhinus mrigala and *Oreochromis niloticus*. LAP LAMBERT Academic Publishing GmbH & Co. KG, Dudweiler Landstr. 99, 66123

- Saarbrücken, Germany 188.
38. Abouelfadl KY, Aly W, Osman AGM (2020) Ageing Nile tilapia (*Oreochromis niloticus*): A comparative study between scales and otoliths. *Int. J. Aquat. Biol* 8: 262-271.
 39. Mayank P, Dwivedi AC, Pathak RK (2018) Age, growth and age pyramid of exotic fish species *Oreochromis niloticus* (Linnaeus 1758) from the lower stretch of the Yamuna river, India. *National Academy Science Letter* 41: 345-348.
 40. El-Kasheif MA, Authman MMN, Al-Ghamdi FA, Ibrahim SA, El-Far AM (2015) Biological aspects and fisheries management of Tilapia fish *Oreochromis niloticus* (Linnaeus, 1758) in El-Bahr El-Faraouny canal, Al-Minufiya Province, Egypt. *Journal of Fisheries and Aquatic Science* 10: 405-444.
 41. Dwivedi AC (2006) Age structure of some commercially exploited fish stocks of the Ganga river system (Banda-Mirzapur section). Thesis submitted to Department of Zoology, University of Allahabad, Prayagraj, (Uttar Pradesh) 138.
 42. Hatem MH, Marwa MM (2008) Biology and fisheries management of Tilapia species in Rosetta branch of the Nile river, Egypt. *Egypt J Aqu Res* 34: 272-284.
 43. Arlinghaus R, Mastsumura S, Dieckmann U (2010) The conservation and fishery benefits of protecting large pike (*Exos lucius* L.) by harvesting regulations in recreational fishing. *Biological Cons* 143: 1444-1459.
 44. Lowe MR, Wu W, Peterson MS, Brown-Peterson NJ, Slack WT, (2012) Survival, growth and reproduction of non-native Nile Tilapia II: fundamental niche projections and invasion potential in the Northern Gulf of Mexico. *PLoS ONE* 7.
 45. Dwivedi AC, Mayank P, Tiwari A (2017) Size selectivity of active fishing gear: changes in size, age and growth of *Cirrhinus mrigala* from the Ganga River, India. *Fisheries and Aquaculture Journal* 8.
 46. Pathak RK, Gopesh A, Dwivedi AC, Joshi KD (2014) Age and growth of alien fish species, *Cyprinus carpio* var. *communis* (Common carp) in the lower stretch of the Yamuna river at Allahabad. *National Academy of Science Letter* 37: 419-422.
 47. Prakash S, Mayank P, Dwivedi AC, Mishra BK, Tiwari A, Upadhyay SK (2019) Size structure and exploitation pattern of Rita rita (Hamilton 1822) in the Yamuna river at Prayagraj, Uttar Pradesh. *Journal of the Kalash Science* 7: 21-24.
 48. Dwivedi AC, Mayank P, Tripathi S (2011) Size composition, exploitation structure and sex ratio of catfish, Rita rita (Hamilton) in the lower stretch of the Yamuna river at Allahabad. *Flora and Fauna* 17: 295-300.
 49. Dwivedi AC, Mayank P, Pathak RK (2016) Size composition and exploitation structure of Indian major carp, *Cirrhinus mrigala* (Hamilton, 1822) from the Ganga river, India. *Journal of Fisheries and Life Science* 1: 30-32.
 50. Imran S, Thakur S, Jha DN, Dwivedi AC (2015) Size composition and exploitation pattern of *Labeo calbasu* (Hamilton 1822) from the lower stretch of the Yamuna river. *Asian Journal of Bio Science* 10: 162-164.
 51. Khan S, Dwivedi AC, Mayank P (2017) Size composition and exploitation pattern of *Cirrhinus mrigala* (Hamilton, 1822) from the Ghaghara River, India. *Journal of Scientific Achievements* 2: 20-22.
 52. Mayank P, Dwivedi AC (2016) Stock assessment and population structure of alien fish species, *Oreochromis niloticus* (Linnaeus) from the lower stretch of the Yamuna river, India. *Journal of the Experimental Zoology, India*, 19: 163-167.
 53. Nautiyal P, Dwivedi AC (2019) Fishery in the tributaries of Yamuna river (Ken river, Paisuni rivers) and Ganga river (Tons river). *Journal of Mountain Research* 14: 19-36.
 54. Dwivedi AC, Mayank P, Shadab Masud S, Khan S (2009) An investigation of the population status and age pyramid of *Cyprinus carpio* var. *communis* from the Yamuna river at Allahabad. *The Asian Journal of Animal Science* 4: 98-101.
 55. Dwivedi AC, Jha DN, Das SCS, Mayank P (2017) Population structure of Nile tilapia, *Oreochromis niloticus* (Linnaeus 1758) from the Ken River, India. *Journal of Scientific Achievements* 2: 23-27.
 56. Mayank P, Rizvi AF, Dwivedi AC (2017) Population dynamics of *Cirrhinus mrigala* (Hamilton 1822) from the largest tributary of the Ganga River, India. *International Journal of Fauna and Biological Studies* 4: 42-47.
 57. Gopesh A, Tripathi S, Joshi KD, Dwivedi AC (2020) Size composition, exploitation structure and sex ratio of *Clupisoma garua* (Hamilton) from middle stretch of the Ganga River at Allahabad, India. *National Academy Science Letter*, <https://doi.org/10.1007/s40009-020-01011-0>.
 58. Toussaint A, Beauchard O, Oberdorff T, Brosse S, Villeger S (2016) Worldwide freshwater fish homogenization is driven by a few widespread non-native species. *Biological Invasions* 18:1295.
 59. Dwivedi, AC, Tiwari A, Mayank P (2018) Environmental pollution supports to constancy and invader potential of *Cyprinus carpio* and *Oreochromis niloticus* from the Ganga river, India. *International Journal of Poultry and Fisheries Sciences* 2: 1-7.
 60. Tiwari A, Dwivedi AC (2014) Assessment of heavy metals bioaccumulation in alien fish species *Cyprinus carpio* from the Gomti river, India. *European Journal of Experimental Biology* 4: 112-117.
 61. Tiwari A, Dwivedi AC, Rahman MA (2020) Assessment of heavy metal accumulation in vital tissues of commercially exploited fish, *Cyprinus carpio* (common carp) from the river Ganga, India. *Journal of Emerging Technologies and Innovative Research* 7: 959-972.
 62. Dwivedi AC, Tiwari A, Mayank P (2015) Seasonal determination of heavy metals in muscle, gill and liver tissues of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) from the tributary of the Ganga River, India. *Zoology and Ecology* 25: 166-171.
 63. Harting JH, Zarull MA, Ciborowski J, Gannon J, Wilke E, et al. (2009) Long-term ecosystem monitoring and assessment of the Detroit river and western lake Erie. *Environ. Monit. Assess* 158: 87-104.
 64. Hering D, Aroviita J, Baattrup, Pedersen A (2015) contrasting the roles of section length and instream habitat enhancement for river restoration success: a field study of 20 European restoration projects. *J. Appl. Ecol* doi: 10.1111/1365-2664.12531.
 65. AC, Nautiyal P (2010) Population dynamics of important fishes in the Vindhyan region, India. LAP LAMBERT Academic Publishing GmbH & Co. KG, Dudweiler Landstr 99, 66123 Saarbrücken, Germany 220.

Copyright: ©2021 Amitabh Chandra Dwivedi. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.