

## Research Article

## Open Access

## Pattern and Outcome of Dog Bite Injuries among Children-a Study in 200 Cases

Waliur Rahman<sup>1\*</sup>, Habibullah Sarkar<sup>2</sup>, Samir Kumar Talukder<sup>3</sup>, Joynal Abedin<sup>4</sup> and Paritosh Kumar Ghosh<sup>5</sup>

<sup>1</sup>Senior Consultant (Surgery), Chuadanga Sadar Hospital, Chuadanga, Bangladesh

<sup>2</sup>Professor of Surgery, Rajshahi Medical College, Rajshahi, Bangladesh

<sup>3</sup>Associate professor (Endocrinology), Rangpur Medical College, Rangpur, Bangladesh

<sup>4</sup>Assistant professor (Surgery), Jashore Medical College, Jashore, Bangladesh

<sup>5</sup>Assistant professor (Cardiology), Kushtia Medical College, Kushtia, Bangladesh

### ABSTRACT

**Introduction:** Dog bites in humans are a major public health problem. Globally, millions of people are bitten by dogs but most of the fatal cases occur in children. Dog bites in human are a serious public health problem and have been well documented worldwide. As rabies is not a notifiable disease in Bangladesh and most deaths occur in rural areas where surveillance is poor.

**Objectives:** To determine the pattern of dog bite injuries and associated health problems among children.

**Methods:** The study was an observational retrospective study carried out at the Dept. of General Surgery, Chuadanga Sadar Hospital, Chuadanga, Bangladesh. The study reviewed the clinical data of patients managed for dog bite related injuries and rabies over a four and half year period between January 2016 and June 2020. A proforma was designed to extract relevant clinical data from the case records. Information extracted included the age, sex of the victims, site of the bite, time of presentation in the hospital, pre-hospital treatment, hospital treatment including post-exposure prophylaxis and complication.

**Results:** In all, 200 cases of dog bite injuries were managed constituting 0.89% of the total consultations; 5 (2.5%) had rabies. Most of the victims were aged 6-12 years (55.0%) and majority (67.0%) was boys. Eighty two percent of the victims presented within 24hrs of the injury. 92 (46.0%) had WHO grade 3 dog bite injury at presentation and the lower limb was the commonest (56.0%) bite site. Use of herbal preparation was the most common pre-hospital treatment 60%. Although 95.0% received anti-rabies vaccine, only 55.5% of them completed the vaccination schedule. The case fatality rate for dog bite was 5.0%. The 5 that died all presented late, had no post exposure prophylaxis and died within 24 hours of admission.

**Conclusion:** There is need for public enlightenment on dangers associated with dog bites and also for the government to ensure vaccination for cost of post exposure prophylaxis treatment for children free of cost.

### \*Corresponding author

Md.Waliur Rahman, Senior Consultant, Chuadanga Sadar Hospital, Chuadanga, Bangladesh; Email: publication985@gmail.com

**Received:** August 21, 2021; **Accepted:** August 26, 2021; **Published:** August 31, 2021

**Keywords:** Rabies, Prescriptions, Rabies Vaccine, Rabies Immunoglobulin.

### Introduction

Dog bites in human are a serious public health problem and have been well documented worldwide [1, 2]. In the United States, 4.7 million people were estimated to have been bitten by dogs in 1994 (an incidence rate of 16.1/1000 in adults and 24.5/1000 in children), of whom 800,000 required medical treatment [3]. The annual estimated number of dog bites in India is 17.4 million, leading to estimated 18,000-20,000 cases of human rabies per

year [4]. Rabies, which is a 100% fatal and 100% preventable disease, mainly occur by dog bite. Dog bite remains a major public health and clinical problem causing palpable fear and anxiety in the patients, the relatives and the attending health care worker not only for the associated morbidity but for the risk of contracting rabies which if not adequately prevented or passes unrecognized almost always result in fatal outcome [5-7]. Dogs are the major reservoir and vector of the rabies virus in Bangladesh [8]. Dog bites and rabies are a major public health issue globally and this is not unconnected with the well-known fact that rabies is usually fatal once the clinical symptoms develop [9]. Dog bites are the

main causes of these preventable traumatic injuries especially in the Paediatric population [10]. Globally, millions of people are bitten by dogs; reports in the USA estimated about 4.5 million dog bites per year and this resulted in 368,265 emergency department visits in 2001 [11]. It was reported that, forty-two percent of dog bite victims are children below fourteen years of age and the injury rate was reported to be highest for children between 5 and 9 years of age with the rate decreasing with increasing age [11]. The injury rate was also reported to be higher among the boys than girls [11]. Dog bites cause a crushing-type wound because of their rounded teeth and strong jaws. Dog bites can result in varying degrees of injuries but fatal injuries occur mostly in children. Victims suffer from psychological and emotional trauma in addition to the physical injuries sustained from the bites [10]. Stray and unvaccinated dogs abound on the streets in Bangladesh and other developing countries thereby putting children particularly at risk of either provoked or unprovoked attack by these dogs. Many of these dogs are dogs initially kept as pets, for security purpose, for hunting and also for consumption in some cultures; many of them are later abandoned by the owners. Almost all of them have never had rabies vaccination thus putting victims at risk of rabies [5, 6]. The most feared complication of dog bites is rabies although not all bites result in rabies [12]. Rabies is an acute viral disease of the central nervous system that remains largely untreatable. Rabies (meaning madness in Latin) is caused by rabies virus of the family Rhabdoviridae and the genus *Lyssavirus* [13]. The rabies virus affects all warm-blooded animals including man and the outcome is usually fatal [14]. It causes an acute encephalomyelitis and it is typically transmitted from the saliva of a rabid animal via a bite, scratch or mucous membrane exposure [9]. Dogs and cats are the major reservoirs and vectors of rabies transmission especially in Africa and Asia whereas in the Americas, bats are more important in transmitting the disease [15]. Rabies is almost always fatal if post-exposure prophylaxis (PEP) is not administered before the development of symptoms [9]. Globally, about 55,000 human deaths occur annually and over 98 percent of these mortalities are caused by canine rabies [16]. The only way of preventing rabies is through the administration of pre-exposure immunization to individuals at risk, administration of PEP with or without human rabies immune globulin (HRIG) and control of rabies in animal population [17]. Human diploid cell vaccine in combination with rabies immune globulin when administered promptly to rabies-exposed patients after appropriate wound care is virtually 100% effective in preventing human death [17]. This study was carried out to determine the prevalence and pattern as well as the outcome of dog bite injuries among children seen at Secondary Care Hospital, Bangladesh.

## Methods

The study was an observational retrospective study carried out at the Dept. of General Surgery, Chuadanga Sadar Hospital, Chuadanga, Bangladesh. The study reviewed the clinical data of patients managed for dog bite related injuries and rabies over a four and half year period between January 2016 and June 2020. A proforma was designed to extract relevant clinical data from the case records. Information extracted included the age, sex of the victims, site of the bite, time of presentation in the hospital, pre-hospital treatment, hospital treatment including post-exposure prophylaxis and complication. The severity of the injury was determined using the WHO dog bite injury grading system [18].

## Data analysis

The information obtained was entered into personal computer and analysed using SPSS version 18. Data was presented in numbers and percentages, chi square was used to compare groups and a p-value of <0.05 was considered significant.



**Figure 1:** Several dog bite injuries

## Results

A total 200 cases of dog bite injuries with clinical features of rabies. Majority of the patients 164 (82.0%), presented within 24 hours of sustaining the dog bite injury while 24 (12.0%) presented between 24 to 48 hours and the remaining 12 (6.0%) after several weeks. As shown in (Table-1), most of the victims are in the age range 6 – 12 years 110 (55.0%), 134 boys (67.07%) were involved giving a male: female ratio of 2.0:1. About 92.5% of the affected children were from the middle and lower socioeconomic classes. 81 children (40.5%) presented with wounds while 20 (10.0%) victims presented for the first time, in the hospital with central nervous system symptoms weeks after sustaining dog bite injury. The lower limb was the most common site of bite in about 56% of cases (Table-2). 92 (46.0%) of the victims presented with grade III wound severity grading as shown in (Figure-2). Human rabies immune globulin was administered to all patients' grade III injury. Although 182 (91.0%) of the victims had post exposure prophylaxis (PEP) with anti-rabies vaccine (Table 1), only 100 (50.0%) of them were managed successfully and subsequently discharged after ensuring adequate wound healing and completion of the vaccination regimen. 81 (40.5%) were lost to follow up and did not complete the vaccination series, the remaining 5 victims (2.5%) all, from the lower socioeconomic class who presented late with features of rabies such as hydrophobia, restlessness and aggression died within 24 hours of admission.

The treatment administered is determined by the severity or grade of the injury. Patients with higher grades of injury significantly received more hospital treatments ( $P=0.001$ ) (Table 3). Human rabies immune globulin was not administered on any of the patients. Although 182 (91.0%) of the victims had post-exposure prophylaxis (PEP) with anti-rabies vaccine (Table 1), only 100 (50.0%) of them were managed successfully and subsequently discharged after ensuring adequate wound healing and completion of the vaccination regimen. 81 (40.5%) were lost to follow up and did not complete the vaccination series, the remaining 12 victims (6.0%) all, from the lower socioeconomic class who presented late with features of rabies such as hydrophobia, restlessness and aggression died within 24 hours of admission. All the patients who died presented weeks after sustaining dog bite injuries and they did not receive any anti rabies vaccination before presentation thereby giving rabies a case fatality rate of 100% in this study. None among those who had anti-rabies vaccine died ( $P=0.001$ ) (Table 4). Post mortem examination was not done on any of the victims who died because the parents or care givers declined consent for post mortem examination.


**Table 1: Socio-demographic characteristics and symptoms of dog bite victims (N=200)**

Variable	Frequency	Percentage (%)
Age range		
0 – 5 yrs	50	25.0
6 – 12 yrs	110	55.0
> 12 yrs	40	20.0
Total	200	100
Sex		
Male	134	67.00
Female	66	33.00
Total	200	100
Social Class		
Lower	111	55.5
Middle	84	42.00

Upper	25	12.5
Total	200	100
Presenting symptom		
Wounds	161	82.0
Lacerations	18	9.00
Bleeding	36	18.00
Fever	12	6.00
Pain/ Swellings	24	12.00
CNS symptoms	12	6.00

**Table 2: Sites of dog bite injury (N=200)**

Site of injury	Number of victims	Percentage (%)
Lower limb	112	56.0
Upper limb	50	25.0
Head & Neck	20	10.00
Trunk/ Buttocks	18	9.00
Total	200	100



**Figure 2: Dog bite wound severity grading in the patients**

**Table 3: Comparison of injury grade with treatment received at the hospital (N=200)**

		Injury grade			Total
		Grade-1	Grade-2	Grade-3	
Treatment at Hospital	Anti-Rabies & Tetanus prophylaxis	0	28	45	77
	Anti-Rabies & Wound dressing	9	15	15	33
	Anti-Rabies & Analgesics	0	6	14	23
	Wound dressing, Antibiotics & Tetanus prophylaxis	0	0	9	9
	Wound dressing, Antibiotics, Tetanus prophylaxis & Analgesics	0	6	0	6
Total		27	81	92	200
p-value = 0.001					

Treatments received in the hospital ranged from washing the bite site with soap and water, to suturing of lacerations and wound dressing, analgesics, tetanus prophylaxis, anti-rabies vaccination (ARV), intravenous fluids and diazepam administration as well as antibiotics administration. The treatment administered is determined by the severity or grade of the injury. Patients with higher grades of injury significantly received more hospital treatments ( $P=0.001$ ) (Table-3).



**Table 4: Outcome of treatment (N=200)**

		Outcome			Total
		Dead	Discharged	Lost to Follow up	
ARV Administration	No	10	0	0	10
	Yes	0	110	80	190
Total		10	110	80	200

p-value = 0.001; NB: ARV-Anti Rabies Vaccine

All the patients who died presented weeks after sustaining dog bite injuries and they did not receive any anti rabies vaccination before presentation thereby giving rabies a case fatality rate of 100% in this study. None among those who had anti-rabies vaccine died (p=0.001) Post mortem examination was not done on any of the victims who died because the parents or care givers declined consent for post mortem examination (Table-4).

## Discussion

This study showed that dog bite related injuries are not a common presentation both in the children emergency unit as well as the out-patient department of our hospital as it constituted about 0.89% of the total consultation over the study period. This figure is however higher than 0.31% and 0.24% reported by Schalamon et al and Kahn et al in their studies respectively [19, 20]. This prevalence is however less than 1.5% documented by Dwyer et al at a pediatric trauma unit in South Africa [21]. For the families that lost their wards to dog bite related injuries, it was a 100% loss hence, dog bites still remains a major public health issue in most developing countries because these deaths can be prevented. The high vulnerability of children to dog bite injury as shown by previous studies conducted locally and internationally has been attributed to children's inability to defend themselves, their small stature and also to the fact that they may provoke the attack in the first instance [8, 10, 21, 22]. Whereas adults are able to stand up to the dogs and defend themselves in most cases, there were many reported cases of unprovoked attacks in this study further highlighting the need to protect children from undue attacks by dogs in the study area. It might be important to come up with prevention programs for children since they are the most vulnerable group as evidenced by this study. Prevention programs such as teaching children how to respond to, behave around and interact with dogs; a randomized controlled trial of a school based intervention in Australia documented a substantial reduction in children's approach to and interaction with a strange dog [23]. This study revealed that children between the ages of 6 and 12 years are more commonly bitten by dogs and also showed male preponderance (71.4%) among the victims which is similar to findings in previous studies and this is not unconnected with the natural behaviour of boys who are more adventurous and daring than their female counterparts, they are also likely to yell, play with or grab the dogs if possible, which put them at risk of being bitten [8, 22, 24, 25]. The most common site of bites in this study was the lower limb which is similar to what was documented by Abubakar et al in Jos and Iyalohme et al in Ekpoma, Edo state. Similar to these studies, head and neck injuries were uncommon in our study [5, 8 & 4]. Majority of the bites were from stray dogs whose owners were known to the victims. This is a common practice in our environment as dog owners mostly do not restrain their dogs and they allow them to wander about the streets. Many of these dogs are not well fed by their owners and as such wander around looking for food. No wonder that there was evidence of vaccination in just one (1.2%) of the cases seen in our hospital

during the study period. Hence, mass vaccination of domestic dogs as well as removal of stray dogs can help in reducing the risk of dog bites as this has successfully eliminated or controlled domestic dog rabies in many parts of the world [26]. Responsible dog ownership should also be included as part of the preventive measures although this seems not to be feasible with the current economic situation in most homes in the country. Post exposure administration of human diploid cell vaccine in combination with human rabies immune globulin soon after bite by suspected rabid dogs or other canines have been found to be very effective in controlling human rabies [16]. The recommended post exposure prophylaxis is intramuscular administration of human diploid cell vaccine into the deltoid on days 0, 3, 7, 14, and 28. Injections can be given into anterolateral thigh for children [27, 28]. Only a little above 50% of the dog bite victims in this study completed the PEP regimen which is slightly lower than the 60.5% documented by Abubakar et al in Jos [5]. This is probably not unconnected with the cost of the vaccine as most of the victims in this study were from the middle and lower socioeconomic classes. A single vial of anti-rabies vaccine (ARV costs 500 Tk., HRIG 1000 Tk.) alone besides other adjunct materials needed for its administration. Given the high preponderance of poverty in Bangladesh, it may be important for the government to look at the possibility of providing this vaccine free of cost [8]. The development of a safe, inexpensive anti-rabies vaccine that can afford adequate protection from a single dose will also go a long way in protecting the huge population of children who are at risk. All the patients with grade III injury vaccine HRIG. This may also be due to the fact that the attending physicians did not prescribe it alongside the human diploid cell vaccine or that the care givers couldn't afford it because of the exorbitant cost or probably because it was not available. Eighty-two percent of the victims in this study presented within 24 hours of injury which is commendable but this is slightly lower than 87.7% reported by Abubakar et al in Jos [5]. Most of our patients had inappropriate home remedies administered before presentation in the hospital and this could have delayed hospital presentation in some of them thus putting the victims at risk of developing complications or even a false sense of assurance when in the real sense the problem was yet to fully evolve. Hence, public enlightenment campaign would be of help in encouraging early presentation as well as informing the public on the right thing to do if they encounter such victims. The five deaths recorded in this study, were from the lower social class, they did not receive PEP with ARV and also presented late with features of rabies and all of them died within 24 hours of presentation. These observations not only underscore the importance of timely PEP in preventing rabies following dog bite injury, but also highlight the unholy alliances between poverty, ignorance, and inadequate access to quality health care in preventing avoidable deaths. The parents of the affected children might have acted out of ignorance or lack of financial power to be able to access care and purchase the human diploid vaccine to complete the PEP regimen among other treatments. There is therefore the need for more aggressive health education of the public and also for the provision of subsidy on the vaccine including the human rabies immune globulin as well as other cares for children with dog bite injury. This will make health care affordable at our health facilities and help to prevent avoidable deaths.

## Conclusion

In conclusion, this study revealed that dog bite related injuries and rabies are not common reasons for consultation at the children emergency and pediatric outpatient units. However, it highlights the need for public health awareness on the health challenges posed

by dog bites because of its high case fatality rate as shown in this study. It also points out the need for public health education on the immediate management of dog bite victims and the preventive measures that need to be put in place to prevent dog bite injuries and also inappropriate pre-hospital management as well as ensuring the availability and affordability of the human diploid cell vaccine and the human rabies immune globulin at our public health facilities. Caregivers should seek medical care in earnest whenever their wards are involved in dog bite injury as delaying may lead to grievous health consequences.

## References

- Overall KL, Love M (2001) Dog bites to humans: demography, epidemiology, injury, and risk. *J Am Vet Med Assoc* 218: 1923-1934.
- Ozanne-Smith J, Ashby K, Stathakis VZ (2001) Dog bite and injury prevention-analysis, critical review, and research agenda 7: 321-326.
- Sacks JJ, Kresnow M, Houston B (1996) Dog bites: How big a problem? 2: 52-54.
- Gongal G, Wright AE (2011) Human rabies in the WHO Southeast Asia region: Forward steps for elimination Article ID 383870.
- Abubakar SA, Bakari AG (2012) Incidence of dog bite injuries and clinical rabies in a tertiary health care institution: a 10-year retrospective study 11: 108-111.
- Bata SI, Ozikwi A, Ayika DG (2011) Retrospective study of dog bite cases reported to ECWA Veterinary Clinic, Bukuru, and Plateau State 6: 17-19.
- Adeleke SI (2010) Impact of dog bite in Kano City: a retrospective study 13: 67-69.
- M. A. H. N. A. Khan (2015) Important Vector-Borne Diseases with Their Zoonotic Potential: Present Situation and Future Perspective, *Bangl. J. Vet. Med* 13: 1-14.
- Iyalomhe GBS, Iyalomhe SI (2014) Dog bite and clinical rabies in a suburban hospital in Nigeria: a 20-year retrospective study of the prevalence and treatment with anti-rabies vaccine. *World J Pharm Res* 4: 113-121.
- Dwyer JP, Douglas TS, van As AB (2007) Dog bite injuries in children - a review of data from a South African pediatric trauma unit. *S Afr Med J*. 2007; 97(8): 597-600.
- CDC. Nonfatal injuries treated in hospital emergency departments-United States, 2001. *MMWR Morb Mortal Wkly Rep* 52: 605-610.
- Alabi O, Nguku P, Chukwukere S, Gaddo A, Nsubuga P, et.al.(2014) Profile of dog bite victims in Jos Plateau State, Nigeria: a review of dog bite records (2006-2008). *Pan Afr Med J* 18: 12.
- Simpson DP (1979) *Cassell's Latin Dictionary*. 5th Ed. London, Cassell Ltd.
- Adeyemi I, Zessin K (2000) Retrospective dog rabies vaccination evaluation at the University Of Ibadan, Nigeria (1988 – 1992). *Veterinarski ARHIV* 70: 223-230.
- Wunner WH, Briggs DJ (2010) Rabies in the 21st century. *PLoS Negl Trop Dis* 4: e591.
- Aworh MK, Nwosuh CI, Ajumobi OO, Okewole PA, Okolocha EC, et.al. (2011) A retrospective study of rabies cases reported at Vom Christian hospital, Plateau State, Nigeria, 2006 - 2010. *Nig Veter J* 32: 366-370.
- Uzzaman MS (2011) Rabies: taking notice of neglected disease Accessed September 24th.
- WHO (2004) Report of the WHO expert consultation on rabies. Geneva: WHO.
- Schalamon J, Ainoedhofer H, Singer G, Petnehazy T, Mayr J et.al. (2006) Analysis of dog bites in children who are younger than 17yrs. *Pediatrics* 117: e374- e379.
- Kahn A, Bauche P, Lamoureux J (2003) Child victims of dog bites treated in emergency departments: a prospective survey. *Eur J Pediatr* 162: 254-258.
- Dwyer JP, Douglas TS, van As AB (2007) Dog bite injuries in children - a review of data from a South African pediatric trauma unit. *S Afr Med J* 97: 597-600.
- Ahmed H, Chafe UM, Magaji AA, Abdul-Qadir A (2000) Rabies and dog bite in children: A decade of experience in Sokoto, Nigeria. *Sokoto J Vet Sci* 1: 2-10.
- Aghahowa SE, Ogbevoen RN (2010) Incidence of dog bite and anti- rabies vaccine utilization in the University of Benin Teaching Hospital, Benin City, Nigeria: A 12-year assessment. *Vaccine* 28: 4847-4850.
- Chapman S, Cornwall J, Righetti J, Sung L (2000) Preventing dog bites in children: randomised controlled trial of an educational intervention 320: 1512-1513.
- Hampson K, Dushoff J, Bingham J, Brückner G, Ali YH, et.al (2007) Synchronous cycles of domestic dogs rabies in sub-Saharan Africa and the impact of control efforts. *Proc Natl Acad Sci USA* 104 : 7717-7722.
- Warell MJ (2000) Rabies encephalitis and its prophylaxis. *Pract Neurol*. 2001; 1: 14-29.
- Ahmed H, Chafe UM, Magaji AA, Abdul-Qadir A. Rabies and dog bite in children: A decade of experience in Sokoto, Nigeria. *Sokoto J Vet Sci* 1: 2-10.
- Aghahowa SE, Ogbevoen RN (2010) Incidence of dog bite and anti- rabies vaccine utilization in the University of Benin Teaching Hospital, Benin City, Nigeria: A 12-year assessment. *Vaccine* 28: 4847-4850.

**Copyright:** ©2021 Waliur Rahman, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.